

CHAPTER 11

Limits and an Introduction to Calculus

Section 11.1	Introduction to Limits	929
Section 11.2	Techniques for Evaluating Limits	936
Section 11.3	The Tangent Line Problem	947
Section 11.4	Limits at Infinity and Limits of Sequences	959
Section 11.5	The Area Problem	966
Review Exercises	976
Practice Test	989

CHAPTER 11

Limits and an Introduction to Calculus

Section 11.1 Introduction to Limits

- If $f(x)$ becomes arbitrarily close to a unique number L as x approaches c from either side, then the limit of $f(x)$ as x approaches c is L :

$$\lim_{x \rightarrow c} f(x) = L.$$

- You should be able to use a calculator to find a limit.
- You should be able to use a graph to find a limit.
- You should understand how limits can fail to exist:
 - $f(x)$ approaches a different number from the right of c than it approaches from the left of c .
 - $f(x)$ increases or decreases without bound as x approaches c .
 - $f(x)$ oscillates between two fixed values as x approaches c .
- You should know and be able to use the elementary properties of limits.


Vocabulary Check

1. limit


2. oscillates

3. direct substitution

1. (a)



(b) $V = (\text{base})\text{height} = (24 - 2x)^2x = 4x(12 - x)^2$




(c) $\lim_{x \rightarrow 4} V = 1024$

Maximum at $x = 4$


x	3	3.5	3.9	4	4.1	4.5	5
V	972.0	1011.5	1023.5	1024.0	1023.5	1012.5	980.0

2. (a)



(b) $x^2 + y^2 = 18 \Rightarrow y = \sqrt{18 - x^2}$

Area = $\frac{1}{2}bh = \frac{1}{2}x\sqrt{18 - x^2}$



(c) $\lim_{x \rightarrow 3} A(x) = 4.5$

x	2	2.5	2.9	3	3.1	3.5	4
A	3.74	4.28	4.49	4.5	4.49	4.20	2.83

3. $\lim_{x \rightarrow 2} (5x + 4) = 14$

x	1.9	1.99	1.999	2	2.001	2.01	2.1
$f(x)$	13.5	13.95	13.995	14	14.005	14.05	14.5

The limit is reached.

4. $\lim_{x \rightarrow 1} (2x^2 + x - 4) = -1$

x	0.9	0.99	0.999	1	1.001	1.01	1.1
$f(x)$	-1.48	-1.0498	-1.0050	-1	-0.9950	-0.9498	-0.48

The limit is reached.

5. $\lim_{x \rightarrow 3} \frac{x - 3}{x^2 - 9} = \frac{1}{6}$

x	2.9	2.99	2.999	3	3.001	3.01	3.1
$f(x)$	0.1695	0.1669	0.16669	Error	0.16664	0.1664	0.1639

The limit is not reached.

6. $\lim_{x \rightarrow -1} \frac{x + 1}{x^2 - x - 2} = -\frac{1}{3}$

x	-1.1	-1.01	-1.001	-1.0	-0.999	-0.99	-0.9
$f(x)$	-0.3226	-0.3322	-0.3332	Error	-0.3334	-0.3344	-0.3348

The limit is not reached.

7. $\lim_{x \rightarrow 0} \frac{\sin 2x}{x} = 2$

x	-0.1	-0.01	-0.001	0	0.001	0.01	0.1
$f(x)$	1.9867	1.99987	1.9999987	Error	1.9999987	1.99987	1.987

The limit is not reached.

8. $\lim_{x \rightarrow 0} \frac{\tan x}{2x} = \frac{1}{2}$

x	-0.1	-0.01	-0.001	0	0.001	0.01	0.1
$f(x)$	0.5017	0.50002	0.5	Error	0.5	0.50002	0.5017

The limit is not reached.

$$9. \lim_{x \rightarrow 0} \frac{e^{2x} - 1}{x} = 2$$

x	-0.1	-0.01	-0.001	0	0.001	0.01	0.1
$f(x)$	1.8127	1.9801	1.9980	Error	2.0020	2.0201	2.2140

The limit is not reached.


$$10. \lim_{x \rightarrow 1} \frac{\ln x}{x - 1} = 1$$

x	0.9	0.99	0.999	1	1.001	1.01	1.1
$f(x)$	1.0536	1.0050	1.0005	Error	0.9995	0.9950	0.9531

The limit is not reached.


$$11. \lim_{x \rightarrow 1} \frac{x - 1}{x^2 + 2x - 3} = \frac{1}{4}$$

x	0.9	0.99	0.999	1.0	1.001	1.01	1.1
$f(x)$	0.2564	0.2506	0.2501	Error	0.2499	0.2494	0.2439




$$12. \lim_{x \rightarrow -2} \frac{x + 2}{x^2 + 5x + 6} = 1$$

x	-2.1	-2.01	-2.001	-2.0	-1.999	-1.99	-1.9
$f(x)$	1.1111	1.0101	1.0010	Error	0.9990	0.9901	0.9091




$$13. \lim_{x \rightarrow 0} \frac{\sqrt{x+5} - \sqrt{5}}{x} \approx 0.2236 \quad \left(\text{Actual limit is } \frac{1}{2\sqrt{5}} \right)$$

x	-0.1	-0.01	-0.001	0	0.001	0.01	0.1
$f(x)$	0.2247	0.2237	0.2236	Error	0.2236	0.2235	0.2225




$$14. \lim_{x \rightarrow -3} \frac{\sqrt{1-x} - 2}{x + 3} = -\frac{1}{4}$$

x	-3.1	-3.01	-3.001	-3.0	-2.999	-2.99	-2.9
$f(x)$	-0.2485	-0.2498	-0.25	Error	-0.25	-0.2502	-0.2516




$$15. \lim_{x \rightarrow -4} \frac{[x/(x+2)] - 2}{x + 4} = \frac{1}{2}$$

x	-4.1	-4.01	-4.001	-4.0	-3.999	-3.99	-3.9
$f(x)$	0.4762	0.4975	0.4998	Error	0.5003	0.5025	0.5263



$$16. \lim_{x \rightarrow 2} \frac{\frac{1}{x+2} - \frac{1}{4}}{x-2} = -\frac{1}{16}$$


x	1.9	1.99	1.999	2.0	2.001	2.01	2.1
$f(x)$	-0.0641	-0.0627	-0.0625	Error	-0.0625	-0.0623	-0.0610



17. Make sure your calculator is set in radian mode.


$$\lim_{x \rightarrow 0} \frac{\sin x}{x} = 1$$

x	-0.1	-0.01	-0.001	0	0.001	0.01	0.1
$f(x)$	0.9983	0.99998	0.9999998	Error	0.9999998	0.99998	0.9983



$$18. \lim_{x \rightarrow 0} \frac{\cos x - 1}{x} = 0$$

x	-0.1	-0.01	-0.001	0	0.001	0.01	0.1
$f(x)$	0.050	0.005	0.0005	Error	-0.0005	-0.005	-0.050



$$19. \lim_{x \rightarrow 0} \frac{\sin^2 x}{x} = 0$$

x	-0.1	-0.01	-0.001	0	0.001	0.01	0.1
$f(x)$	-0.0997	-0.0100	-0.0010	Error	0.0010	0.0100	0.0997

$$20. \lim_{x \rightarrow 0} \frac{2x}{\tan 4x} = 0.5$$

x	-0.1	-0.01	-0.001	0	0.001	0.01	0.1
$f(x)$	0.4730	0.4997	0.499997	Error	0.499997	0.4997	0.4730

$$21. \lim_{x \rightarrow 0} \frac{e^{2x} - 1}{2x} = 1.0$$

x	-0.1	-0.01	-0.001	0	0.001	0.01	0.1
$f(x)$	0.9063	0.9901	0.9990	Error	1.0010	1.0101	1.1070

$$22. \lim_{x \rightarrow 0} \frac{1 - e^{-4x}}{x} = 4.0$$

x	-0.1	-0.01	-0.001	0	0.001	0.01	0.1
$f(x)$	4.9182	4.0811	4.0080	Error	3.9920	3.9211	3.2968

23. $\lim_{x \rightarrow 1} \frac{\ln(2x - 1)}{x - 1} = 2$

x	0.9	0.99	0.999	1	1.001	1.01	1.1
$f(x)$	2.2314	2.0203	2.0020	Error	1.9980	1.9803	1.8232


24. $\lim_{x \rightarrow 1} \frac{\ln x^2}{x - 1} = 2.0$

x	0.9	0.99	0.999	1	1.001	1.01	1.1
$f(x)$	2.1072	2.0101	2.0010	Error	1.9990	1.9901	1.9062


25. $f(x) = \begin{cases} 2x + 1, & x < 2 \\ x + 3, & x \geq 2 \end{cases}$

The limit exists as x approaches 2:


$$\lim_{x \rightarrow 2} f(x) = 5$$




26. $\lim_{x \rightarrow 2} f(x) = 4$



27. $\lim_{x \rightarrow 2} f(x)$ does not exist.



28. $\lim_{x \rightarrow 2} f(x)$ does not exist.



29. $\lim_{x \rightarrow -4} (x^2 - 3) = 13$

30. $\lim_{x \rightarrow 2} \frac{3x^2 - 12}{x - 2} = 12$

31. $\lim_{x \rightarrow -2} \frac{|x + 2|}{x + 2}$ does not exist. $f(x) = \frac{|x + 2|}{x + 2}$ equals -1 to the left of -2 , and equals 1 to the right of -2 .

32. The limit does not exist because $f(x)$ does not approach a real number as x approaches 1.


33. The limit does not exist because $f(x)$ oscillates between 2 and -2 .

34. $\lim_{x \rightarrow -1} \sin\left(\frac{\pi x}{2}\right) = -1$


35. $\lim_{x \rightarrow \pi/2} \tan x$ does not exist.

36. $\lim_{x \rightarrow \pi/2} \sec x$ does not exist.

37. $\lim_{x \rightarrow 0} \frac{5}{2 + e^{1/x}}$ does not exist.




38. $\lim_{x \rightarrow 0} \frac{e^x - 1}{x} = 1$




39. $\lim_{x \rightarrow 0} \cos \frac{1}{x}$ does not exist.


The graph oscillates between -1 and 1 .




40. $\lim_{x \rightarrow -1} \sin \pi x = 0$




41. $\lim_{x \rightarrow 4} \frac{\sqrt{x+3} - 1}{x-4}$ does not exist.




42. $\lim_{x \rightarrow 2} \frac{\sqrt{x+5} - 4}{x-2}$ does not exist.




43. $\lim_{x \rightarrow 1} \frac{x-1}{x^2 - 4x + 3} = -\frac{1}{2}$




44. $\lim_{x \rightarrow 3} \frac{7}{x-3}$ does not exist.



45. $\lim_{x \rightarrow 4} \ln(x+3) \approx 1.946$ (Exact limit is $\ln 7$.)



46. $\lim_{x \rightarrow -1} \ln(7-x) = \ln(7 - (-1)) = \ln 8$



47. (a) $\lim_{x \rightarrow c} [-2g(x)] = -2(6) = -12$
 (b) $\lim_{x \rightarrow c} [f(x) + g(x)] = 3 + 6 = 9$
 (c) $\lim_{x \rightarrow c} \frac{f(x)}{g(x)} = \frac{3}{6} = \frac{1}{2}$
 (d) $\lim_{x \rightarrow c} \sqrt{f(x)} = \sqrt{3}$

48. (a) $\lim_{x \rightarrow c} [f(x) + g(x)]^2 = (5 - 2)^2 = 9$
 (b) $\lim_{x \rightarrow c} [6f(x)g(x)] = 6(5)(-2) = -60$
 (c) $\lim_{x \rightarrow c} \frac{5g(x)}{4f(x)} = \frac{5(-2)}{4(5)} = -\frac{1}{2}$
 (d) $\lim_{x \rightarrow c} \frac{1}{\sqrt{f(x)}} = \frac{1}{\sqrt{5}} = \frac{\sqrt{5}}{5}$

49. (a) $\lim_{x \rightarrow 2} f(x) = 2^3 = 8$
 (b) $\lim_{x \rightarrow 2} g(x) = \frac{\sqrt{2^2 + 5}}{2(2^2)} = \frac{3}{8}$

- (c) $\lim_{x \rightarrow 2} [f(x)g(x)] = 8\left(\frac{3}{8}\right) = 3$
 (d) $\lim_{x \rightarrow 2} [g(x) - f(x)] = \frac{3}{8} - 8 = -\frac{61}{8}$

50. (a) $\lim_{x \rightarrow 2} f(x) = \frac{2}{3-2} = 2$
 (b) $\lim_{x \rightarrow 2} g(x) = \sin(\pi 2) = 0$
 (c) $\lim_{x \rightarrow 2} [f(x)g(x)] = 2(0) = 0$
 (d) $\lim_{x \rightarrow 2} [g(x) - f(x)] = 0 - 2 = -2$
51. $\lim_{x \rightarrow 5} (10 - x^2) = 10 - 5^2 = -15$
52. $\lim_{x \rightarrow -2} \left(\frac{1}{2}x^3 - 5x \right) = \frac{1}{2}(-2)^3 - 5(-2) = 6$
53. $\lim_{x \rightarrow -3} (2x^2 + 4x + 1) = 2(-3)^2 + 4(-3) + 1 = 7$
54. $\lim_{x \rightarrow -2} (x^3 - 6x + 5) = (-2)^3 - 6(-2) + 5 = 9$
55. $\lim_{x \rightarrow 3} \left(-\frac{9}{x} \right) = -\frac{9}{3} = -3$
56. $\lim_{x \rightarrow -5} \frac{6}{x+2} = \frac{6}{-5+2} = \frac{6}{-3} = -2$
57. $\lim_{x \rightarrow -3} \frac{3x}{x^2+1} = \frac{9}{10}$
58. $\lim_{x \rightarrow 4} \frac{x-1}{x^2+2x+3} = \frac{4-1}{16+8+3} = \frac{3}{27} = \frac{1}{9}$
59. $\lim_{x \rightarrow -2} \frac{5x+3}{2x-9} = \frac{5(-2)+3}{2(-2)-9} = \frac{-7}{-13} = \frac{7}{13}$
60. $\lim_{x \rightarrow 3} \frac{x^2+1}{x} = \frac{9+1}{3} = \frac{10}{3}$
61. $\lim_{x \rightarrow -1} \sqrt{x+2} = \sqrt{-1+2} = 1$
62. $\lim_{x \rightarrow 3} \sqrt[3]{x^2-1} = \sqrt[3]{9-1} = 2$
63. $\lim_{x \rightarrow 7} \frac{5x}{\sqrt{x+2}} = \frac{5(7)}{\sqrt{7+2}} = \frac{35}{3}$
64. $\lim_{x \rightarrow 8} \frac{\sqrt{x+1}}{x-4} = \frac{\sqrt{8+1}}{8-4} = \frac{3}{4}$
65. $\lim_{x \rightarrow 3} e^x = e^3 \approx 20.0855$
66. $\lim_{x \rightarrow e} \ln x = \ln e = 1$
67. $\lim_{x \rightarrow \pi} \sin 2x = \sin 2\pi = 0$
68. $\lim_{x \rightarrow \pi} \tan x = \tan \pi = 0$
69. $\lim_{x \rightarrow 1/2} \arcsin x = \arcsin \frac{1}{2} = \frac{\pi}{6} \approx 0.5236$
70. $\lim_{x \rightarrow 1} \arccos \frac{x}{2} = \arccos \frac{1}{2} = \frac{\pi}{3} \approx 1.0472$
71. True
72. True (assuming the limits exist).
73. Answers will vary.
74. In general, you cannot use a graphing utility to determine whether a limit can be reached. It is important to analyze a function analytically.
75. (a) No. The limit may or may not exist, and if it does exist, it may not equal 4.
 (b) No. $f(2)$ may or may not exist, and if $f(2)$ exists, it may not equal 4.
76. $\lim_{x \rightarrow 5} f(x) = 12$ means that the values of f approach 12 as x approaches 5.

$$77. \frac{5-x}{3x-15} = \frac{5-x}{-3(5-x)} = -\frac{1}{3}, x \neq 5$$

$$78. \frac{x^2-81}{9-x} = \frac{(x-9)(x+9)}{9-x} = -x-9, x \neq 9$$

$$79. \frac{15x^2+7x-4}{15x^2+x-2} = \frac{(3x-1)(5x+4)}{(3x-1)(5x+2)}$$

$$= \frac{5x+4}{5x+2}, x \neq \frac{1}{3}$$

$$80. \frac{x^2-12x+36}{x^2-7x+6} = \frac{(x-6)(x-6)}{(x-6)(x-1)} = \frac{x-6}{x-1}, x \neq 6$$

$$81. \frac{x^2+27}{x^2+x-6} = \frac{(x+3)(x^2-3x+9)}{(x+3)(x-2)}$$

$$= \frac{x^2-3x+9}{x-2}, x \neq -3$$

$$82. \frac{x^3-8}{x^2-4} = \frac{(x-2)(x^2+2x+4)}{(x-2)(x+2)}$$

$$= \frac{x^2+2x+4}{x+2}, x \neq 2$$

Section 11.2 Techniques for Evaluating Limits

- You can use direct substitution to find the limit of a polynomial function $p(x)$:

$$\lim_{x \rightarrow c} p(x) = p(c).$$

- You can use direct substitution to find the limit of a rational function $r(x) = \frac{p(x)}{q(x)}$, as long as $q(c) \neq 0$:

$$\lim_{x \rightarrow c} r(x) = r(c) = \frac{p(c)}{q(c)}, q(c) \neq 0.$$

- You should be able to use cancellation techniques to find a limit.
- You should know how to use rationalization techniques to find a limit.
- You should know how to use technology to find a limit.
- You should be able to calculate one-sided limits.

Vocabulary Check

1. dividing out technique 2. indeterminate form 3. one-sided limit 4. difference quotient

1. $g(x) = \frac{-2x^2+x}{x}$, $g_2(x) = -2x+1$

(a) $\lim_{x \rightarrow 0} g(x) = 1$

(b) $\lim_{x \rightarrow -1} g(x) = 3$

(c) $\lim_{x \rightarrow -2} g(x) = 5$

2. $h(x) = \frac{x^2-3x}{x}$, $h_2(x) = x-3$

(a) $\lim_{x \rightarrow -2} h(x) = -5$

(b) $\lim_{x \rightarrow 0} h(x) = -3$

(c) $\lim_{x \rightarrow 3} h(x) = 0$

$$3. g(x) = \frac{x^3 - x}{x - 1}, g_2(x) = x^2 + x = x(x + 1)$$

$$(a) \lim_{x \rightarrow 1} g(x) = 2$$

$$(b) \lim_{x \rightarrow -1} g(x) = 0$$

$$(c) \lim_{x \rightarrow 0} g(x) = 0$$

$$4. f(x) = \frac{x^2 - 1}{x + 1}, f_2(x) = x - 1$$

$$(a) \lim_{x \rightarrow 1} f(x) = 0$$

$$(b) \lim_{x \rightarrow 2} f(x) = 1$$

$$(c) \lim_{x \rightarrow -1} f(x) = -2$$

$$5. \lim_{x \rightarrow 6} \frac{x - 6}{x^2 - 36} = \lim_{x \rightarrow 6} \frac{x - 6}{(x - 6)(x + 6)}$$

$$= \lim_{x \rightarrow 6} \frac{1}{x + 6} = \frac{1}{12}$$

$$6. \lim_{x \rightarrow 9} \frac{9 - x}{x^2 - 81} = \lim_{x \rightarrow 9} \frac{9 - x}{(x - 9)(x + 9)}$$

$$= \lim_{x \rightarrow 9} \frac{-1}{x + 9} = \frac{-1}{18}$$

$$7. \lim_{x \rightarrow -1} \frac{1 - 2x - 3x^2}{1 + x} = \lim_{x \rightarrow -1} \frac{(1 + x)(1 - 3x)}{1 + x}$$

$$= \lim_{x \rightarrow -1} (1 - 3x) = 4$$

$$8. \lim_{x \rightarrow -4} \frac{2x^2 + 7x - 4}{x + 4} = \lim_{x \rightarrow -4} \frac{(x + 4)(2x - 1)}{x + 4}$$

$$= \lim_{x \rightarrow -4} (2x - 1) = -9$$

$$9. \lim_{t \rightarrow 2} \frac{t^3 - 8}{t - 2} = \lim_{t \rightarrow 2} \frac{(t - 2)(t^2 + 2t + 4)}{t - 2}$$

$$= \lim_{t \rightarrow 2} (t^2 + 2t + 4)$$

$$= 4 + 4 + 4 = 12$$

$$10. \lim_{a \rightarrow -4} \frac{a^3 + 64}{a + 4} = \lim_{a \rightarrow -4} \frac{(a + 4)(a^2 - 4a + 16)}{a + 4}$$

$$= \lim_{a \rightarrow -4} (a^2 - 4a + 16)$$

$$= 16 + 16 + 16 = 48$$

$$11. \lim_{x \rightarrow 1} \frac{x^4 - 1}{x^4 - 3x^2 - 4} = \frac{0}{-6} = 0$$

$$12. \lim_{x \rightarrow 2} \frac{x^4 - 2x^2 - 8}{x^4 - 6x^2 + 8} = \lim_{x \rightarrow 2} \frac{(x - 2)(x + 2)(x^2 + 2)}{(x - 2)(x + 2)(x^2 - 2)}$$

$$= \lim_{x \rightarrow 2} \frac{(x + 2)(x^2 + 2)}{(x + 2)(x^2 - 2)}$$

$$= \frac{4(6)}{4(2)} = 3$$

$$13. \lim_{x \rightarrow -1} \frac{x^3 + 2x^2 - x - 2}{x^3 + 4x^2 - x - 4} = \lim_{x \rightarrow -1} \frac{(x - 1)(x + 1)(x + 2)}{(x - 1)(x + 1)(x + 4)}$$

$$= \lim_{x \rightarrow -1} \frac{(x - 1)(x + 2)}{(x - 1)(x + 4)}$$

$$= \frac{(-2)(1)}{(-2)(3)} = \frac{1}{3}$$

$$14. \lim_{x \rightarrow -3} \frac{x^3 + 2x^2 - 9x - 18}{x^3 + x^2 - 9x - 9} = \lim_{x \rightarrow -3} \frac{(x - 3)(x + 2)(x + 3)}{(x - 3)(x + 1)(x + 3)}$$

$$= \lim_{x \rightarrow -3} \frac{x + 2}{x + 1}$$

$$= \frac{-1}{-2} = \frac{1}{2}$$

$$\begin{aligned}
 15. \lim_{x \rightarrow 2} \frac{x^3 + 2x^2 - 5x - 6}{x^3 - 7x + 6} &= \lim_{x \rightarrow 2} \frac{(x-2)(x+1)(x+3)}{(x-2)(x-1)(x+3)} \\
 &= \lim_{x \rightarrow 2} \frac{x+1}{x-1} \\
 &= \frac{3}{1} = 3
 \end{aligned}$$

$$\begin{aligned}
 16. \lim_{x \rightarrow 3} \frac{x^3 - 4x^2 - 3x + 18}{x^3 - 4x^2 + x + 6} &= \lim_{x \rightarrow 3} \frac{(x-3)^2(x+2)}{(x-3)(x-2)(x+1)} \\
 &= \lim_{x \rightarrow 3} \frac{(x-3)(x+2)}{(x-2)(x+1)} \\
 &= \frac{0}{4} = 0
 \end{aligned}$$

$$\begin{aligned}
 17. \lim_{y \rightarrow 0} \frac{\sqrt{5+y} - \sqrt{5}}{y} &= \lim_{y \rightarrow 0} \frac{\sqrt{5+y} - \sqrt{5}}{y} \cdot \frac{\sqrt{5+y} + \sqrt{5}}{\sqrt{5+y} + \sqrt{5}} \\
 &= \lim_{y \rightarrow 0} \frac{(5+y) - 5}{y(\sqrt{5+y} + \sqrt{5})} \\
 &= \lim_{y \rightarrow 0} \frac{1}{\sqrt{5+y} + \sqrt{5}} \\
 &= \frac{1}{2\sqrt{5}} = \frac{\sqrt{5}}{10}
 \end{aligned}$$

$$\begin{aligned}
 18. \lim_{z \rightarrow 0} \frac{\sqrt{7-z} - \sqrt{7}}{z} \left(\frac{\sqrt{7-z} + \sqrt{7}}{\sqrt{7-z} + \sqrt{7}} \right) &= \lim_{z \rightarrow 0} \frac{(7-z) - 7}{z(\sqrt{7-z} + \sqrt{7})} \\
 &= \lim_{z \rightarrow 0} \frac{-1}{\sqrt{7-z} + \sqrt{7}} \\
 &= \frac{-1}{2\sqrt{7}} = -\frac{\sqrt{7}}{14}
 \end{aligned}$$

$$\begin{aligned}
 19. \lim_{x \rightarrow -3} \frac{\sqrt{x+7} - 2}{x+3} &= \lim_{x \rightarrow -3} \frac{\sqrt{x+7} - 2}{x+3} \cdot \frac{\sqrt{x+7} + 2}{\sqrt{x+7} + 2} \\
 &= \lim_{x \rightarrow -3} \frac{(x+7) - 4}{(x+3)(\sqrt{x+7} + 2)} \\
 &= \lim_{x \rightarrow -3} \frac{1}{\sqrt{x+7} + 2} = \frac{1}{4}
 \end{aligned}$$

$$\begin{aligned}
 20. \lim_{x \rightarrow 2} \frac{4 - \sqrt{18-x}}{x-2} &= \lim_{x \rightarrow 2} \frac{4 - \sqrt{18-x}}{x-2} \cdot \frac{4 + \sqrt{18-x}}{4 + \sqrt{18-x}} \\
 &= \lim_{x \rightarrow 2} \frac{16 - (18-x)}{(x-2)(4 + \sqrt{18-x})} \\
 &= \lim_{x \rightarrow 2} \frac{1}{4 + \sqrt{18-x}} = \frac{1}{8}
 \end{aligned}$$


$$\begin{aligned}
 21. \lim_{x \rightarrow 0} \frac{1/(1+x) - 1}{x} &= \lim_{x \rightarrow 0} \frac{1 - (1+x)}{(1+x)x} \\
 &= \lim_{x \rightarrow 0} \frac{-1}{1+x} = -1
 \end{aligned}$$

$$\begin{aligned}
 23. \lim_{x \rightarrow 0} \frac{\sec x}{\tan x} &= \lim_{x \rightarrow 0} \frac{1}{\cos x} \cdot \frac{\cos x}{\sin x} \\
 &= \lim_{x \rightarrow 0} \frac{1}{\sin x}, \text{ does not exist}
 \end{aligned}$$

$$\begin{aligned}
 25. \lim_{x \rightarrow 0} \frac{\cos 2x}{\cot 2x} &= \lim_{x \rightarrow 0} \frac{\cos 2x}{(\cos 2x)/\sin(2x)} \\
 &= \lim_{x \rightarrow 0} \sin 2x = 0
 \end{aligned}$$

$$27. \lim_{x \rightarrow \pi/2} \frac{\sin x - 1}{x} = \frac{1 - 1}{\pi/2} = 0$$

$$\begin{aligned}
 29. f(x) &= \frac{\sqrt{x+3} - \sqrt{3}}{x} \\
 \lim_{x \rightarrow 0} f(x) &\approx 0.2887 \\
 \left(\text{Exact limit: } \frac{1}{2\sqrt{3}} \right)
 \end{aligned}$$




$$\begin{aligned}
 22. \lim_{x \rightarrow 0} \frac{\frac{1}{x-8} + \frac{1}{8}}{x} &= \lim_{x \rightarrow 0} \frac{8 + (x-8)}{(x-8)(8)x} \\
 &= \lim_{x \rightarrow 0} \frac{1}{(x-8)8} = -\frac{1}{64}
 \end{aligned}$$

$$\begin{aligned}
 24. \lim_{x \rightarrow \pi/2} \frac{1 - \sin x}{\cos x} &= \lim_{x \rightarrow \pi/2} \frac{1 - \sin x}{\cos x} \cdot \frac{1 + \sin x}{1 + \sin x} \\
 &= \lim_{x \rightarrow \pi/2} \frac{1 - \sin^2 x}{\cos x(1 + \sin x)} \\
 &= \lim_{x \rightarrow \pi/2} \frac{\cos^2 x}{\cos x(1 + \sin x)} \\
 &= \lim_{x \rightarrow \pi/2} \frac{\cos x}{1 + \sin x} = 0
 \end{aligned}$$


$$\begin{aligned}
 26. \lim_{x \rightarrow 0} \frac{\sin x - x}{\sin x} &= \lim_{x \rightarrow 0} \left(1 - \frac{x}{\sin x} \right) \\
 &= 1 - 1 = 0
 \end{aligned}$$

$$28. \lim_{x \rightarrow \pi} \frac{1 + \cos x}{x} = \frac{1 + (-1)}{\pi} = 0$$


$$30. \lim_{x \rightarrow 0} \frac{\sqrt{x+4} - 2}{x} \approx 0.25, \left(\frac{1}{4} \right)$$




$$\begin{aligned}
 31. f(x) &= \frac{\sqrt{2x+1} - 1}{x} \\
 \lim_{x \rightarrow 0} f(x) &= 1
 \end{aligned}$$




$$32. \lim_{x \rightarrow 9} \frac{3 - \sqrt{x}}{x - 9} = -0.1667, \left(-\frac{1}{6} \right)$$



$$33. \lim_{x \rightarrow 2} \frac{x^5 - 32}{x - 2} = 80$$




$$34. \lim_{x \rightarrow 1} \frac{x^4 - 1}{x - 1} = 4$$




35. $f(x) = \frac{1/(x+4) - (1/4)}{x}$

$\lim_{x \rightarrow 0} f(x) = -\frac{1}{16}, (-0.0625)$




36. $\lim_{x \rightarrow 0} \frac{\frac{1}{2+x} - \frac{1}{2}}{x} = -0.25, \left(-\frac{1}{4}\right)$




37. $f(x) = \frac{e^{2x} - 1}{x}$


$\lim_{x \rightarrow 0} f(x) = 2$




38. $\lim_{x \rightarrow 0} \frac{1 - e^{-x}}{x} = 1$




39. $\lim_{x \rightarrow 0^+} x \ln x = 0$




40. $\lim_{x \rightarrow 0^+} x^2 \ln x = 0$




41. $\lim_{x \rightarrow 0} \frac{\sin 2x}{x} = 2$




42. $\lim_{x \rightarrow 0} \frac{\sin 3x}{x} = 3$




43. $\lim_{x \rightarrow 0} \frac{\tan x}{x} = 1$




44. $\lim_{x \rightarrow 0} \frac{1 - \cos 2x}{x} = 0$



45. $\lim_{x \rightarrow 1} \frac{1 - \sqrt[3]{x}}{1 - x} = \frac{1}{3} \approx 0.333$




46. $\lim_{x \rightarrow 1} \frac{\sqrt[3]{x} - x}{x - 1} \approx -0.667, \left(-\frac{2}{3}\right)$




47. $f(x) = (1 - x)^{2/x}$

$\lim_{x \rightarrow 0} f(x) \approx 0.135$




48. $\lim_{x \rightarrow 0} (1 + 2x)^{1/x} \approx 7.389$



$$49. f(x) = \frac{x-1}{x^2-1}$$

(a) Graphically, $\lim_{x \rightarrow 1^-} \frac{x-1}{x^2-1} = \frac{1}{2}$.




(b)

x	0.5	0.9	0.99	0.999	1
$f(x)$	0.6667	0.5263	0.5025	0.5003	Error

Numerically, $\lim_{x \rightarrow 1^-} \frac{x-1}{x^2-1} = \frac{1}{2}$.

(c) Algebraically, $\lim_{x \rightarrow 1^-} \frac{x-1}{x^2-1} = \lim_{x \rightarrow 1^-} \frac{x-1}{(x-1)(x+1)} = \lim_{x \rightarrow 1^-} \frac{1}{x+1} = \frac{1}{2}$.

$$50. \lim_{x \rightarrow 5^+} \frac{5-x}{25-x^2} = 0.1$$



(b)


x	5.1	5.01	5.001	5
$f(x)$	0.099	0.0999	0.09999	Error

(c) Algebraically,

$$\begin{aligned} \lim_{x \rightarrow 5^+} \frac{5-x}{25-x^2} &= \lim_{x \rightarrow 5^+} \frac{(5-x)}{(5-x)(5+x)} \\ &= \lim_{x \rightarrow 5^+} \frac{1}{5+x} = \frac{1}{10} \end{aligned}$$

$$51. f(x) = \frac{4-\sqrt{x}}{x-16}$$

(a) Graphically, $\lim_{x \rightarrow 16^+} \frac{4-\sqrt{x}}{x-16} = -\frac{1}{8}$.



(b)


x	16	16.001	16.01	16.1	16.5
$f(x)$	Error	-0.1250	-0.1250	-0.1248	-0.1240

Numerically, $\lim_{x \rightarrow 16^+} \frac{4-\sqrt{x}}{x-16} = -0.125$.

(c) Algebraically,

$$\begin{aligned} \lim_{x \rightarrow 16^+} \frac{4-\sqrt{x}}{x-16} &= \lim_{x \rightarrow 16^+} \frac{4-\sqrt{x}}{(\sqrt{x}-4)(\sqrt{x}+4)} \\ &= \lim_{x \rightarrow 16^+} \frac{-1}{\sqrt{x}+4} = \frac{-1}{4+4} = -\frac{1}{8} \end{aligned}$$

52. $\lim_{x \rightarrow 0^-} \frac{\sqrt{x+2} - \sqrt{2}}{x} \approx 0.3536$



(b)

x	-1.0	-0.1	-0.01	-0.001	0
$f(x)$	0.4142	0.3581	0.3540	0.3536	?

(c) Algebraically,


$$\begin{aligned} \lim_{x \rightarrow 0^-} \frac{\sqrt{x+2} - \sqrt{2}}{x} \cdot \frac{\sqrt{x+2} + \sqrt{2}}{\sqrt{x+2} + \sqrt{2}} &= \lim_{x \rightarrow 0^-} \frac{(x+2) - 2}{x(\sqrt{x+2} + \sqrt{2})} \\ &= \lim_{x \rightarrow 0^-} \frac{1}{\sqrt{x+2} + \sqrt{2}} = \frac{1}{2\sqrt{2}} = \frac{\sqrt{2}}{4} \approx 0.3536. \end{aligned}$$

53. $f(x) = \frac{|x-6|}{x-6}$

$\lim_{x \rightarrow 6^+} f(x) = 1$

$\lim_{x \rightarrow 6^-} f(x) = -1$

Limit does not exist.




54. $\lim_{x \rightarrow 2^-} \frac{|x-2|}{x-2} = -1$

$\lim_{x \rightarrow 2^+} \frac{|x-2|}{x-2} = 1$

$\lim_{x \rightarrow 2} \frac{|x-2|}{x-2} = 1$

Limit does not exist.




55. $f(x) = \frac{1}{x^2+1}$

$\lim_{x \rightarrow 1^-} \frac{1}{x^2+1} = \lim_{x \rightarrow 1^+} \frac{1}{x^2+1}$

$= \lim_{x \rightarrow 1} \frac{1}{x^2+1}$


$= \frac{1}{2}$



56. $\lim_{x \rightarrow 1^-} \frac{1}{x^2-1}$ does not exist.

$\lim_{x \rightarrow 1^+} \frac{1}{x^2-1}$ does not exist.


$\lim_{x \rightarrow 1} \frac{1}{x^2-1}$ does not exist.



57. $\lim_{x \rightarrow 2^-} f(x) = 2 - 1 = 1$

$\lim_{x \rightarrow 2^+} f(x) = 2(2) - 3 = 1$


$\lim_{x \rightarrow 2} f(x) = 1$



58. $\lim_{x \rightarrow 1^-} f(x) = 2(1) + 1 = 3$

$\lim_{x \rightarrow 1^+} f(x) = 4 - 1 = 3$

$\lim_{x \rightarrow 1} f(x) = 3$




$$59. f(x) = \begin{cases} 4 - x^2, & x \leq 1 \\ 3 - x, & x > 1 \end{cases}$$

$$\lim_{x \rightarrow 1^-} f(x) = 4 - 1 = 3$$

$$\lim_{x \rightarrow 1^+} f(x) = 3 - 1 = 2$$



$$\lim_{x \rightarrow 1} f(x) \text{ does not exist.}$$




$$60. \lim_{x \rightarrow 0^-} f(x) = 4 - 0 = 4$$

$$\lim_{x \rightarrow 0^+} f(x) = 0 + 4 = 4$$


$$\lim_{x \rightarrow 0} f(x) = 4$$




$$\lim_{x \rightarrow 0} f(x) = 0$$




$$\lim_{x \rightarrow 0} f(x) = 0$$




$$\lim_{x \rightarrow 0} f(x) = 0$$



$$\lim_{x \rightarrow 0} f(x) = 0$$



$$\lim_{x \rightarrow 0} f(x) = 0$$



$$\lim_{x \rightarrow 0} f(x) = 0$$

67. (a) Can be evaluated by direct substitution:

$$\lim_{x \rightarrow 0} x^2 \sin x^2 = 0^2 \sin 0^2 = 0$$

(b) Cannot be evaluated by direct substitution:

$$\lim_{x \rightarrow 0} \frac{\sin x^2}{x^2} = 1$$

68. (a) Can be evaluated by direct substitution.

$$\lim_{x \rightarrow 0} \frac{x}{\cos x} = \frac{0}{\cos 0} = 0$$

(b) Cannot be evaluated by direct substitution.

$$\lim_{x \rightarrow 0} \frac{1 - \cos x}{x} = 0$$

(See Section 11.1, Exercise 18.)

$$\begin{aligned} 69. \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h} &= \lim_{h \rightarrow 0} \frac{3(x+h) - 1 - (3x - 1)}{h} \\ &= \lim_{h \rightarrow 0} \frac{3x + 3h - 1 - 3x + 1}{h} \\ &= \lim_{h \rightarrow 0} \frac{3h}{h} = 3 \end{aligned}$$

$$\begin{aligned} 70. \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h} &= \lim_{h \rightarrow 0} \frac{[5 - 6(x+h)] - (5 - 6x)}{h} \\ &= \lim_{h \rightarrow 0} -\frac{6h}{h} = -6 \end{aligned}$$

$$\begin{aligned}
 71. \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h} &= \lim_{h \rightarrow 0} \frac{\sqrt{x+h} - \sqrt{x}}{h} \cdot \left(\frac{\sqrt{x+h} + \sqrt{x}}{\sqrt{x+h} + \sqrt{x}} \right) \\
 &= \lim_{h \rightarrow 0} \frac{(x+h) - x}{h(\sqrt{x+h} + \sqrt{x})} \\
 &= \lim_{h \rightarrow 0} \frac{1}{\sqrt{x+h} + \sqrt{x}} = \frac{1}{2\sqrt{x}}
 \end{aligned}$$

$$\begin{aligned}
 72. \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h} &= \lim_{h \rightarrow 0} \frac{\sqrt{x+h-2} - \sqrt{x-2}}{h} \cdot \frac{\sqrt{x+h-2} + \sqrt{x-2}}{\sqrt{x+h-2} + \sqrt{x-2}} \\
 &= \lim_{h \rightarrow 0} \frac{(x+h-2) - (x-2)}{h[\sqrt{x+h-2} + \sqrt{x-2}]} \\
 &= \lim_{h \rightarrow 0} \frac{1}{\sqrt{x+h-2} + \sqrt{x-2}} \\
 &= \frac{1}{2\sqrt{x-2}}
 \end{aligned}$$

$$\begin{aligned}
 73. \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h} &= \lim_{h \rightarrow 0} \frac{((x+h)^2 - 3(x+h)) - (x^2 - 3x)}{h} \\
 &= \lim_{h \rightarrow 0} \frac{x^2 + 2xh + h^2 - 3x - 3h - x^2 + 3x}{h} \\
 &= \lim_{h \rightarrow 0} \frac{2xh + h^2 - 3h}{h} \\
 &= \lim_{h \rightarrow 0} (2x + h - 3) = 2x - 3
 \end{aligned}$$

$$\begin{aligned}
 74. \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h} &= \lim_{h \rightarrow 0} \frac{[4 - 2(x+h) - (x+h)^2] - [4 - 2x - x^2]}{h} \\
 &= \lim_{h \rightarrow 0} \frac{4 - 2x - 2h - x^2 - 2xh - h^2 - 4 + 2x + x^2}{h} \\
 &= \lim_{h \rightarrow 0} \frac{-2h - 2xh - h^2}{h} \\
 &= \lim_{h \rightarrow 0} (-2 - 2x - h) = -2 - 2x
 \end{aligned}$$


$$\begin{aligned}
 75. \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h} &= \lim_{h \rightarrow 0} \frac{1/(x+h+2) - 1/(x+2)}{h} \\
 &= \lim_{h \rightarrow 0} \frac{(x+2) - (x+h+2)}{h(x+h+2)(x+2)} \\
 &= \lim_{h \rightarrow 0} \frac{-h}{h(x+h+2)(x+2)} \\
 &= \lim_{h \rightarrow 0} \frac{-1}{(x+h+2)(x+2)} \\
 &= \frac{-1}{(x+2)^2}
 \end{aligned}$$

$$\begin{aligned}
 76. \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h} &= \lim_{h \rightarrow 0} \frac{\frac{1}{x+h-1} - \frac{1}{x-1}}{h} \\
 &= \lim_{h \rightarrow 0} \frac{x-1 - (x+h-1)}{h(x+h-1)(x-1)} \\
 &= \lim_{h \rightarrow 0} \frac{-1}{(x+h-1)(x-1)} \\
 &= \frac{-1}{(x-1)^2}
 \end{aligned}$$

$$\begin{aligned}
 77. \lim_{t \rightarrow 1} \frac{(-16(1) + 128) - (-16t^2 + 128)}{1-t} &= \lim_{t \rightarrow 1} \frac{16t^2 - 16}{1-t} \\
 &= \lim_{t \rightarrow 1} \frac{16(t-1)(t+1)}{1-t} \\
 &= \lim_{t \rightarrow 1} -16(t+1) \\
 &= -32 \frac{\text{ft}}{\text{sec}}
 \end{aligned}$$

$$\begin{aligned}
 78. v(2) &= \lim_{t \rightarrow 2} \frac{s(2) - s(t)}{2-t} = \lim_{t \rightarrow 2} \frac{(-64 + 128) - (-16t^2 + 128)}{2-t} \\
 &= \lim_{t \rightarrow 2} \frac{16t^2 - 64}{2-t} = \lim_{t \rightarrow 2} \frac{16(t+2)(t-2)}{2-t} \\
 &= \lim_{t \rightarrow 2} -16(t+2) = -64 \text{ feet per second}
 \end{aligned}$$

$$79. C(t) = 1.00 - 0.25[-(t-1)]$$



(b)

t	3	3.3	3.4	3.5	3.6	3.7	4
C	1.50	1.75	1.75	1.75	1.75	1.75	1.75


$$\lim_{t \rightarrow 3.5} C(t) = 1.75$$

(c)

t	2	2.5	2.9	3	3.1	3.5	4
C	1.25	1.50	1.50	1.50	1.75	1.75	1.75

$\lim_{t \rightarrow 3} C(t)$ does not exist. The one-sided limits do not agree.

80. $C(t) = 1.25 - 0.15\lceil -(t - 1) \rceil$



(b)

t	3	3.3	3.4	3.5	3.6	3.7	4
C	1.55	1.70	1.70	1.70	1.70	1.70	1.70

$$\lim_{t \rightarrow 3.5} C(t) = 1.70$$

(c)

t	2	2.5	2.9	3	3.1	3.5	4
C	1.4	1.55	1.55	1.55	1.7	1.7	1.7

$\lim_{t \rightarrow 3} C(t)$ does not exist. The one-sided limits do not agree.



81. Answers will vary. As $t \rightarrow 2$ from the left, $f(t) \rightarrow 39.00$. As $t \rightarrow 2$ from the right, $f(t) \rightarrow 46.80$.

82. Answers will vary. As $x \rightarrow 1$ from the left, $f(x) \rightarrow 14.40$. As $x \rightarrow 1$ from the right, $f(x) \rightarrow 18.30$.

83. True

84. False. The value of f at c has no bearing on the limit.

85. Many answers possible



86. Answers will vary.

87. Slope of line through $(4, -6)$ and $(3, -4)$:

$$\frac{-6 - (-4)}{4 - 3} = -2$$

Slope of perpendicular line: $\frac{1}{2}$

Equation: $y + 10 = \frac{1}{2}(x - 6)$

$$2y - x + 26 = 0$$

88. Slope between $(3, -3)$ and $(5, -2)$:


$$\frac{-2 - (-3)}{5 - 3} = \frac{1}{2}$$

Line: $y + 1 = \frac{1}{2}(x - 1)$


$$2y + 2 = x - 1$$

$$2y - x + 3 = 0$$

$$89. r = \frac{3}{1 + \cos \theta}, e = 1, \text{ Parabola}$$




$$90. r = \frac{12}{3 + 2 \sin \theta} = \frac{4}{1 + (2/3) \sin \theta}, e = \frac{2}{3}, \text{ Ellipse}$$




$$91. r = \frac{9}{2 + 3 \cos \theta} = \frac{9/2}{1 + (3/2) \cos \theta}, e = \frac{3}{2}, \text{ Hyperbola}$$


Hyperbola




$$92. r = \frac{4}{4 + \cos \theta} = \frac{1}{1 + (1/4) \cos \theta}, e = \frac{1}{4}, \text{ Ellipse}$$



$$93. r = \frac{5}{1 - \sin \theta}, e = 1, \text{ Parabola}$$



$$94. r = \frac{6}{3 - 4 \sin \theta} = \frac{2}{1 - (4/3) \sin \theta}, e = \frac{4}{3}, \text{ Hyperbola}$$



$$95. \langle 7, -2, 3 \rangle \cdot \langle -1, 4, 5 \rangle = -7 - 8 + 15 = 0 \Rightarrow \text{orthogonal}$$

$$96. \langle 5, 5, 0 \rangle \cdot \langle 0, 5, 1 \rangle = 25 \neq 0$$

Not multiples of each other; neither parallel nor orthogonal

$$97. -3\langle -4, 3, -6 \rangle = \langle 12, -9, 18 \rangle \Rightarrow \text{parallel}$$

$$98. \langle 2, -3, 1 \rangle \cdot \langle -2, 2, 2 \rangle = -8 \neq 0$$

Not multiples of each other; neither parallel nor orthogonal

Section 11.3 The Tangent Line Problem

- You should be able to visually approximate the slope of a graph.
- The slope m of the graph of f at the point $(x, f(x))$ is given by

$$m = \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h}$$

provided this limit exists.

- You should be able to use the limit definition to find the slope of a graph.
- The derivative of f at x is given by

$$f'(x) = \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h}$$

provided this limit exists. Notice that this is the same limit as that for the tangent line slope.

- You should be able to use the limit definition to find the derivative of a function.

Vocabulary Check

- | | | |
|------------------------|-----------------|----------------|
| 1. Calculus | 2. tangent line | 3. secant line |
| 4. difference quotient | 5. derivative | |

1. Slope is 0 at (x, y) . 2. Slope is -1 at (x, y) . 3. Slope is $\frac{1}{2}$ at (x, y) . 4. Slope is -2 at (x, y) .

$$5. m_{\text{sec}} = \frac{g(3+h) - g(3)}{h} = \frac{(3+h)^2 - 4(3+h) - (-3)}{h} = \frac{h^2 + 2h}{h}$$

$$m = \lim_{h \rightarrow 0} \frac{h^2 + 2h}{h} = \lim_{h \rightarrow 0} \frac{h(h+2)}{h} = \lim_{h \rightarrow 0} (h+2) = 2$$

$$6. m_{\text{sec}} = \frac{f(3+h) - f(3)}{h}$$

$$= \frac{10(3+h) - 2(3+h)^2 - 12}{h}$$

$$= \frac{-2h - 2h^2}{h} = -2 - 2h, h \neq 0$$

$$m = \lim_{h \rightarrow 0} (-2 - 2h) = -2$$

$$7. m_{\text{sec}} = \frac{g(1+h) - g(1)}{h}$$

$$= \frac{5 - 2(1+h) - 3}{h} = \frac{-2h}{h}$$

$$m = \lim_{h \rightarrow 0} \frac{-2h}{h} = -2$$

$$8. m_{\text{sec}} = \frac{h(-1+k) - h(-1)}{k} = \frac{2(-1+k) + 5 - 3}{k} = \frac{2k}{k}$$

$$m = \lim_{k \rightarrow 0} \frac{2k}{k} = 2$$

$$9. m_{\text{sec}} = \frac{g(2+h) - g(2)}{h} = \frac{[4/(2+h)] - 2}{h} = \frac{4 - 2(2+h)}{(2+h)h} = \frac{-2}{2+h}, h \neq 0$$

$$m = \lim_{h \rightarrow 0} \left(\frac{-2}{2+h} \right) = -1$$

$$10. m_{\text{sec}} = \frac{g(4+h) - g(4)}{h} = \frac{\frac{1}{4+h-2} - \frac{1}{2}}{h} = \frac{\frac{1}{2+h} - \frac{1}{2}}{h} = \frac{-h}{(2+h)2h} = \frac{-1}{2(2+h)}, h \neq 0$$

$$m = \lim_{h \rightarrow 0} \left(\frac{-1}{2(2+h)} \right) = -\frac{1}{4}$$

$$11. m_{\text{sec}} = \frac{h(9+k) - h(9)}{k} = \frac{\sqrt{9+k} - 3}{k} \cdot \frac{\sqrt{9+k} + 3}{\sqrt{9+k} + 3} = \frac{(9+k) - 9}{k[\sqrt{9+k} + 3]} = \frac{1}{\sqrt{9+k} + 3}, k \neq 0$$

$$m = \lim_{k \rightarrow 0} \frac{1}{\sqrt{9+k} + 3} = \frac{1}{6}$$

$$12. m_{\text{sec}} = \frac{h(-1+k) - h(-1)}{k} = \frac{\sqrt{-1+k+10} - 3}{k} \cdot \frac{\sqrt{k+9} + 3}{\sqrt{k+9} + 3} = \frac{(k+9) - 9}{k[\sqrt{k+9} + 3]} = \frac{1}{\sqrt{k+9} + 3}, k \neq 0$$

$$m = \lim_{k \rightarrow 0} \frac{1}{\sqrt{k+9} + 3} = \frac{1}{6}$$

$$13. m_{\text{sec}} = \frac{g(x+h) - g(x)}{h} = \frac{4 - (x+h)^2 - (4 - x^2)}{h} = \frac{-2xh - h^2}{h} = -2x - h, h \neq 0$$

$$m = \lim_{h \rightarrow 0} (-2x - h) = -2x$$

$$(a) \text{ At } (0, 4), m = -2(0) = 0.$$

$$(b) \text{ At } (-1, 3), m = -2(-1) = 2.$$

$$14. m_{\text{sec}} = \frac{g(x+h) - g(x)}{h} = \frac{(x+h)^3 - x^3}{h} = \frac{3x^2h + 3xh^2 + h^3}{h}$$

$$= 3x^2 + 3xh + h^2, h \neq 0$$

$$m = \lim_{h \rightarrow 0} (3x^2 + 3xh + h^2) = 3x^2$$

$$(a) \text{ At } (1, 1), m = 3(1)^2 = 3.$$

$$(b) \text{ At } (-2, -8), m = 3(-2)^2 = 12.$$

$$15. m_{\text{sec}} = \frac{g(x+h) - g(x)}{h} = \frac{\frac{1}{x+h+4} - \frac{1}{x+4}}{h} = \frac{(x+4) - (x+4+h)}{(x+h+4)(x+4)(h)}$$

$$= \frac{-h}{(x+h+4)(x+4)h} = \frac{-1}{(x+h+4)(x+4)}, h \neq 0$$

$$m = \lim_{h \rightarrow 0} \frac{-1}{(x+h+4)(x+4)} = \frac{-1}{(x+4)^2}$$

$$(a) \text{ At } \left(0, \frac{1}{4}\right), m = \frac{-1}{(0+4)^2} = \frac{-1}{16}.$$

$$(b) \text{ At } \left(-2, \frac{1}{2}\right), m = \frac{-1}{(-2+4)^2} = \frac{-1}{4}.$$

$$16. m_{\text{sec}} = \frac{f(x+h) - f(x)}{h} = \frac{\frac{1}{x+h+2} - \frac{1}{x+2}}{h} = \frac{(x+2) - (x+h+2)}{h(x+h+2)(x+2)}$$

$$= \frac{-h}{h(x+h+2)(x+2)}$$

$$m = \lim_{h \rightarrow 0} \frac{-h}{h(x+h+2)(x+2)} = \frac{-1}{(x+2)^2}$$

$$(a) \text{ At } \left(0, \frac{1}{2}\right), m = \frac{-1}{(0+2)^2} = \frac{-1}{4}.$$

$$(b) \text{ At } (-1, 1), m = \frac{-1}{(-1+2)^2} = -1.$$

$$17. m_{\text{sec}} = \frac{g(x+h) - g(x)}{h} = \frac{\sqrt{x+h-1} - \sqrt{x-1}}{h} \cdot \frac{\sqrt{x+h-1} + \sqrt{x-1}}{\sqrt{x+h-1} + \sqrt{x-1}}$$

$$= \frac{(x+h-1) - (x-1)}{h(\sqrt{x+h-1} + \sqrt{x-1})} = \frac{1}{\sqrt{x+h-1} + \sqrt{x-1}}, h \neq 0$$

$$m = \lim_{h \rightarrow 0} \left(\frac{1}{\sqrt{x+h-1} + \sqrt{x-1}} \right) = \frac{1}{2\sqrt{x-1}}$$


$$(a) \text{ At } (5, 2), m = \frac{1}{2\sqrt{5-1}} = \frac{1}{4}.$$

$$(b) \text{ At } (10, 3), m = \frac{1}{2\sqrt{10-1}} = \frac{1}{6}.$$


$$\begin{aligned}
 18. m_{\text{sec}} &= \frac{f(x+h) - f(x)}{h} = \frac{\sqrt{x+h-4} - \sqrt{x-4}}{h} \cdot \frac{\sqrt{x+h-4} + \sqrt{x-4}}{\sqrt{x+h-4} + \sqrt{x-4}} \\
 &= \frac{(x+h-4) - (x-4)}{h[\sqrt{x+h-4} + \sqrt{x-4}]} \\
 m &= \lim_{h \rightarrow 0} \frac{h}{h[\sqrt{x+h-4} + \sqrt{x-4}]} = \frac{1}{2\sqrt{x-4}}
 \end{aligned}$$

(a) At (5, 1), $m = \frac{1}{2}$.


(b) At (8, 2), $m = \frac{1}{4}$.




Slope at (1, -1) is 2.




Slope ≈ 0




Slope at (1, 1) is $-\frac{1}{2}$.



At (1, 2), slope $\approx \frac{1}{4}$.



Slope at (1, 2) is -1.



Slope ≈ 3

25. $f'(x) = \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h} = \lim_{h \rightarrow 0} \frac{5 - 5}{h} = 0$

26. $f'(x) = \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h}$
 $= \lim_{h \rightarrow 0} \frac{(-1) - (-1)}{h} = 0$

27. $g'(x) = \lim_{h \rightarrow 0} \frac{g(x+h) - g(x)}{h}$
 $= \lim_{h \rightarrow 0} \frac{[9 - \frac{1}{3}(x+h)] - [9 - \frac{1}{3}x]}{h}$
 $= \lim_{h \rightarrow 0} \frac{-\frac{1}{3}h}{h} = -\frac{1}{3}$

28. $f'(x) = \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h}$
 $= \lim_{h \rightarrow 0} \frac{[-5(x+h) + 2] - (-5x + 2)}{h}$
 $= \lim_{h \rightarrow 0} \frac{-5h}{h} = -5$

29. $f'(x) = \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h} = \lim_{h \rightarrow 0} \frac{[4 - 3(x+h)^2] - (4 - 3x^2)}{h}$
 $= \lim_{h \rightarrow 0} \frac{-3(x^2 + 2xh + h^2) + 3x^2}{h} = \lim_{h \rightarrow 0} \frac{-6xh - 3h^2}{h} = \lim_{h \rightarrow 0} (-6x - 3h) = -6x$

30. $f'(x) = \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h} = \lim_{h \rightarrow 0} \frac{[(x+h)^2 - 3(x+h) + 4] - (x^2 - 3x + 4)}{h}$
 $= \lim_{h \rightarrow 0} \frac{x^2 + 2xh + h^2 - 3h - x^2}{h}$
 $= \lim_{h \rightarrow 0} (2x + h - 3) = 2x - 3$

$$\begin{aligned}
 31. f'(x) &= \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h} \\
 &= \lim_{h \rightarrow 0} \frac{\frac{1}{(x+h)^2} - \frac{1}{x^2}}{h} \\
 &= \lim_{h \rightarrow 0} \frac{x^2 - (x^2 + 2xh + h^2)}{(x+h)^2 x^2 h} \\
 &= \lim_{h \rightarrow 0} \frac{-2x - h}{(x+h)^2 x^2} = -\frac{2x}{x^4} = -\frac{2}{x^3}
 \end{aligned}$$

$$\begin{aligned}
 32. f'(x) &= \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h} = \lim_{h \rightarrow 0} \frac{\frac{1}{(x+h)^3} - \frac{1}{x^3}}{h} \\
 &= \lim_{h \rightarrow 0} \frac{x^3 - (x^3 + 3x^2h + 3xh^2 + h^3)}{h(x+h)^3 x^3} \\
 &= \lim_{h \rightarrow 0} \frac{-3x^2h - 3xh^2 - h^3}{h(x+h)^3 x^3} \\
 &= \lim_{h \rightarrow 0} \frac{-3x^2 - 3xh - h^2}{(x+h)^3 x^3} \\
 &= \frac{-3x^2}{x^6} = \frac{-3}{x^4}
 \end{aligned}$$

$$\begin{aligned}
 33. f'(x) &= \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h} \\
 &= \lim_{h \rightarrow 0} \frac{\sqrt{x+h-4} - \sqrt{x-4}}{h} \cdot \frac{\sqrt{x+h-4} + \sqrt{x-4}}{\sqrt{x+h-4} + \sqrt{x-4}} \\
 &= \lim_{h \rightarrow 0} \frac{(x+h-4) - (x-4)}{h[\sqrt{x+h-4} + \sqrt{x-4}]} \\
 &= \lim_{h \rightarrow 0} \frac{1}{\sqrt{x+h-4} + \sqrt{x-4}} \\
 &= \frac{1}{2\sqrt{x-4}}
 \end{aligned}$$

$$\begin{aligned}
 34. f'(x) &= \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h} \\
 &= \lim_{h \rightarrow 0} \frac{\sqrt{x+h+8} - \sqrt{x+8}}{h} \cdot \frac{\sqrt{x+h+8} + \sqrt{x+8}}{\sqrt{x+h+8} + \sqrt{x+8}} \\
 &= \lim_{h \rightarrow 0} \frac{(x+h+8) - (x+8)}{h[\sqrt{x+h+8} + \sqrt{x+8}]} \\
 &= \lim_{h \rightarrow 0} \frac{1}{\sqrt{x+h+8} + \sqrt{x+8}} \\
 &= \frac{1}{2\sqrt{x+8}}
 \end{aligned}$$

$$\begin{aligned}
 35. f'(x) &= \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h} \\
 &= \lim_{h \rightarrow 0} \frac{\frac{1}{(x+h+2)} - \frac{1}{x+2}}{h} \\
 &= \lim_{h \rightarrow 0} \frac{(x+2) - (x+h+2)}{h(x+h+2)(x+2)} \\
 &= \lim_{h \rightarrow 0} \frac{-1}{(x+h+2)(x+2)} \\
 &= \frac{-1}{(x+2)^2}
 \end{aligned}$$

$$\begin{aligned}
 36. f'(x) &= \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h} \\
 &= \lim_{h \rightarrow 0} \frac{\frac{1}{(x+h-5)} - \frac{1}{x-5}}{h} \\
 &= \lim_{h \rightarrow 0} \frac{(x-5) - (x+h-5)}{h(x+h-5)(x-5)} \\
 &= \lim_{h \rightarrow 0} \frac{-1}{(x+h-5)(x-5)} \\
 &= \frac{-1}{(x-5)^2}
 \end{aligned}$$

$$\begin{aligned}
 37. f'(x) &= \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h} = \lim_{h \rightarrow 0} \frac{\frac{1}{\sqrt{x+h-9}} - \frac{1}{\sqrt{x-9}}}{h} \cdot \frac{\frac{1}{\sqrt{x+h-9}} + \frac{1}{\sqrt{x-9}}}{\frac{1}{\sqrt{x+h-9}} + \frac{1}{\sqrt{x-9}}} \\
 &= \lim_{h \rightarrow 0} \frac{\frac{1}{(x+h-9)} - \frac{1}{(x-9)}}{h \left[\frac{1}{\sqrt{x+h-9}} + \frac{1}{\sqrt{x-9}} \right]} = \lim_{h \rightarrow 0} \frac{(x-9) - (x+h-9)}{h(x+h-9)(x-9) \left[\frac{1}{\sqrt{x+h-9}} + \frac{1}{\sqrt{x-9}} \right]} \\
 &= \lim_{h \rightarrow 0} \frac{-1}{(x+h-9)(x-9) \left[\frac{1}{\sqrt{x+h-9}} + \frac{1}{\sqrt{x-9}} \right]} = \frac{-1}{(x-9)^2 \left[\frac{2}{\sqrt{x-9}} \right]} = \frac{-1}{2(x-9)^{3/2}}
 \end{aligned}$$


$$\begin{aligned}
 38. f'(x) &= \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h} = \lim_{h \rightarrow 0} \frac{\frac{1}{\sqrt{x+h+1}} - \frac{1}{\sqrt{x+1}}}{h} \\
 &= \lim_{h \rightarrow 0} \frac{\sqrt{x+1} - \sqrt{x+h+1}}{h\sqrt{x+h+1}\sqrt{x+1}} \cdot \frac{\sqrt{x+1} + \sqrt{x+h+1}}{\sqrt{x+1} + \sqrt{x+h+1}} \\
 &= \lim_{h \rightarrow 0} \frac{(x+1) - (x+h+1)}{h\sqrt{x+h+1}\sqrt{x+1}[\sqrt{x+1} + \sqrt{x+h+1}]} \\
 &= \lim_{h \rightarrow 0} \frac{-1}{\sqrt{x+h+1}\sqrt{x+1}[\sqrt{x+1} + \sqrt{x+h+1}]} \\
 &= \frac{-1}{\sqrt{x+1}\sqrt{x+1}[2\sqrt{x+1}]} \\
 &= \frac{-1}{2(x+1)^{3/2}}
 \end{aligned}$$

$$\begin{aligned}
 39. (a) m_{\text{sec}} &= \frac{f(2+h) - f(2)}{h} \\
 &= \frac{(2+h)^2 - 1 - 3}{h} \\
 &= \frac{4 + 4h + h^2 - 4}{h} \\
 &= 4 + h, \quad h \neq 0 \\
 m &= \lim_{h \rightarrow 0} (4 + h) = 4
 \end{aligned}$$

$$(b) y - 3 = 4(x - 2)$$

$$y = 4x - 5$$

(c)




$$\begin{aligned}
 40. (a) m_{\text{sec}} &= \frac{f(1+h) - f(1)}{h} \\
 &= \frac{4 - (1+h)^2 - 3}{h} \\
 &= \frac{4 - (1 + 2h + h^2) - 3}{h} \\
 &= \frac{-h^2 - 2h}{h} = -h - 2, \quad h \neq 0 \\
 m &= \lim_{h \rightarrow 0} (-h - 2) = -2
 \end{aligned}$$

$$(b) y - 3 = -2(x - 1)$$

$$y = -2x + 5$$

(c)




$$\begin{aligned}
 41. \quad (a) \quad m_{\text{sec}} &= \frac{f(1+h) - f(1)}{h} \\
 &= \frac{(1+h)^3 - 2(1+h) - (-1)}{h} \\
 &= \frac{1 + 3h + 3h^2 + h^3 - 2 - 2h + 1}{h} \\
 &= \frac{h^3 + 3h^2 + h}{h} = h^2 + 3h + 1, \quad h \neq 0
 \end{aligned}$$

$$m = \lim_{h \rightarrow 0} (h^2 + 3h + 1) = 1$$

$$(b) \quad y - (-1) = 1(x - 1)$$

$$y = x - 2$$

(c)




$$\begin{aligned}
 42. \quad (a) \quad m_{\text{sec}} &= \frac{f(2+h) - f(2)}{h} \\
 &= \frac{(2+h)^3 - (2+h) - 6}{h} \\
 &= \frac{h^3 + 6h^2 + 12h + 8 - 2 - h - 6}{h} \\
 &= \frac{h^3 + 6h^2 + 11h}{h} \\
 &= h^2 + 6h + 11, \quad h \neq 0
 \end{aligned}$$

$$m = \lim_{h \rightarrow 0} (h^2 + 6h + 11) = 11$$

$$(b) \quad y - 6 = 11(x - 2)$$

$$y = 11x - 16$$

(c)




$$\begin{aligned}
 43. \quad (a) \quad m_{\text{sec}} &= \frac{f(3+h) - f(3)}{h} = \frac{\sqrt{3+h+1} - 2}{h} \\
 &= \frac{\sqrt{4+h} - 2}{h} \cdot \frac{\sqrt{4+h} + 2}{\sqrt{4+h} + 2} \\
 &= \frac{4 + h - 4}{h[\sqrt{4+h} + 2]} \\
 &= \frac{1}{\sqrt{4+h} + 2}
 \end{aligned}$$

$$m = \lim_{h \rightarrow 0} \frac{1}{\sqrt{4+h} + 2} = \frac{1}{4}$$

$$(b) \quad y - 2 = \frac{1}{4}(x - 3)$$

$$y = \frac{1}{4}x + \frac{5}{4}$$

(c)




$$\begin{aligned}
 44. \quad (a) \quad m_{\text{sec}} &= \frac{f(3+h) - f(3)}{h} = \frac{\sqrt{3+h-2} - 1}{h} \\
 &= \frac{\sqrt{1+h} - 1}{h} \cdot \frac{\sqrt{1+h} + 1}{\sqrt{1+h} + 1} \\
 &= \frac{(1+h) - 1}{h[\sqrt{1+h} + 1]} \\
 &= \frac{1}{\sqrt{1+h} + 1}, \quad h \neq 0
 \end{aligned}$$

$$m = \lim_{h \rightarrow 0} \frac{1}{\sqrt{1+h} + 1} = \frac{1}{2}$$

$$(b) \quad y - 1 = \frac{1}{2}(x - 3)$$


$$y = \frac{1}{2}x - \frac{1}{2}$$

(c)




45. (a) $m_{\text{sec}} = \frac{f(-4+h) - f(-4)}{h}$
 $= \frac{1}{-4+h+5} - 1$
 $= \frac{1}{h+1} - 1$
 $= \frac{1+h-1-h}{h+1} = \frac{-h}{h+1}, h \neq 0$
 $m = \lim_{h \rightarrow 0} \frac{-h}{h+1} = 0$

(b) $y - 1 = -1(x + 4)$
 $y = -x - 3$




46. (a) $m_{\text{sec}} = \frac{f(4+h) - f(4)}{h}$
 $= \frac{1}{4+h-3} - 1$
 $= \frac{1}{h+1} - 1$
 $= \frac{1-1-h-h}{h+1} = \frac{-2h}{h+1}, h \neq 0$
 $m = \lim_{h \rightarrow 0} \frac{-2h}{h+1} = 0$

(b) $y - 1 = -1(x - 4)$
 $y = -x + 5$



47.

x	-2	-1.5	-1	-0.5	0	0.5	1	1.5	2
$f(x)$	2	1.125	0.5	0.125	0	0.125	0.5	1.125	2
$f'(x)$	-2	-1.5	-1	-0.5	0	0.5	1	1.5	2




$f(x) = \frac{1}{2}x^2$
 $f'(x) = x$

They appear to be the same.

48.

x	-2	-1.5	-1	-0.5	0	0.5	1	1.5	2
$f(x)$	-2	-0.844	-0.25	-0.031	0	0.031	0.25	0.844	2
$f'(x)$	3	1.688	0.75	0.188	0	0.188	0.75	1.688	3




$f(x) = \frac{1}{4}x^3$
 $f'(x) = \frac{3}{4}x^2$

They appear to be the same.

49.

x	-2	-1.5	-1	-0.5	0	0.5	1	1.5	2
$f(x)$	1	1.225	1.414	1.581	1.732	1.871	2	2.121	2.236
$f'(x)$	0.5	0.408	0.354	0.316	0.289	0.267	0.25	0.236	0.224




$$f(x) = \sqrt{x+3}$$

$$f'(x) = \frac{1}{2\sqrt{x+3}}$$

They appear to be the same.

50.

x	-2	-1.5	-1	-0.5	0	0.5	1	1.5	2
$f(x)$	0	-0.7	-1	-1.071	-1	-0.833	-0.6	-0.318	0
$f'(x)$	-2	-0.92	-0.333	0.020	0.25	0.407	0.52	0.603	0.667



$$f(x) = \frac{x^2 - 4}{x + 4}$$

$$f'(x) = \frac{x^2 + 8x + 4}{(x + 4)^2}$$

They appear to be the same.

$$51. f'(x) = \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h} = \lim_{h \rightarrow 0} \frac{[(x+h)^2 - 4(x+h) + 3] - [x^2 - 4x + 3]}{h}$$

$$= \lim_{h \rightarrow 0} \frac{(x^2 + 2xh + h^2 - 4x - 4h + 3) - (x^2 - 4x + 3)}{h}$$

$$= \lim_{h \rightarrow 0} \frac{2xh + h^2 - 4h}{h} = \lim_{h \rightarrow 0} 2x + h - 4 = 2x - 4$$

$$f'(x) = 0 = 2x - 4 \Rightarrow x = 2$$

f has a horizontal tangent at $(2, -1)$.

$$52. f'(x) = \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h}$$

$$= \lim_{h \rightarrow 0} \frac{(x+h)^2 - 6(x+h) + 4 - (x^2 - 6x + 4)}{h}$$

$$= \lim_{h \rightarrow 0} \frac{x^2 + 2xh + h^2 - 6x - 6h - x^2 + 6x}{h}$$

$$= \lim_{h \rightarrow 0} \frac{2xh + h^2 - 6hh}{h} = 2x - 6$$

$$f'(x) = 2x = 6$$

$$f'(x) = 0 \Rightarrow x = 3$$

f has a horizontal tangent at $(3, -5)$.

$$\begin{aligned}
 53. f'(x) &= \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h} = \lim_{h \rightarrow 0} \frac{[3(x+h)^3 - 9(x+h)] - [3x^3 - 9x]}{h} \\
 &= \lim_{h \rightarrow 0} \frac{3(x^3 + 3x^2h + 3xh^2 + h^3) - 9x - 9h - 3x^3 + 9x}{h} \\
 &= \lim_{h \rightarrow 0} \frac{9x^2h + 9xh^2 + 3h^3 - 9h}{h} \\
 &= \lim_{h \rightarrow 0} (9x^2 + 9xh + 3h - 9) = 9x^2 - 9
 \end{aligned}$$

$$f'(x) = 0 = 9x^2 - 9 = 9(x+1)(x-1) \Rightarrow x = \pm 1$$

Horizontal tangents at $(1, -6)$ and $(-1, 6)$

$$\begin{aligned}
 54. f'(x) &= \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h} = \lim_{h \rightarrow 0} \frac{(x+h)^3 + 3(x+h) - (x^3 + 3x)}{h} \\
 &= \lim_{h \rightarrow 0} \frac{x^3 + 3x^2h + 3xh^2 + h^3 + 3x + 3h - x^3 - 3x}{h} \\
 &= \lim_{h \rightarrow 0} \frac{3x^2h + 3xh^2 + h^3 + 3h}{h} \\
 &= \lim_{h \rightarrow 0} (3x^2 + 3xh + h^2 + 3) = 3x^2 + 3
 \end{aligned}$$

$$f'(x) = 3x^2 + 3 = 0, \text{ Impossible}$$

No horizontal tangents

$$\begin{aligned}
 55. \quad f'(x) &= 4x^3 - 4x = 0 \\
 4x(x-1)(x+1) &= 0 \\
 x &= 0, 1, -1 \\
 (0, 0), (1, -1), (-1, -1)
 \end{aligned}$$

$$\begin{aligned}
 56. f'(x) &= 12x^3 + 12x^2 \\
 f'(x) = 0 &= 12x^3 + 12x^2 = 12x^2(x+1) \Rightarrow 0, -1 \\
 f &\text{ has horizontal tangents at } (0, 0) \text{ and } (-1, -1).
 \end{aligned}$$

$$57. f'(x) = -2 \sin x + 1 = 0$$

$$\sin x = \frac{1}{2}$$

$$x = \frac{\pi}{6}, \frac{5\pi}{6}$$

$$\left(\frac{\pi}{6}, \sqrt{3} + \frac{\pi}{6}\right), \left(\frac{5\pi}{6}, \frac{5\pi}{6} - \sqrt{3}\right)$$

$$58. f'(x) = 1 - 2 \cos x = 0$$

$$\cos x = \frac{1}{2}$$

$$x = \frac{\pi}{3}, \frac{5\pi}{3}$$

$$\left(\frac{\pi}{3}, \frac{\pi}{3} - \sqrt{3}\right), \left(\frac{5\pi}{3}, \frac{5\pi}{3} + \sqrt{3}\right)$$

59. $f'(x) = x^2e^x + 2xe^x = 0$

$$xe^x(x + 2) = 0$$

$$x = 0, -2$$

$$(0, 0), (-2, 4e^{-2})$$

60. $f'(x) = e^{-x} - xe^{-x} = 0$

$$e^{-x}(1 - x) = 0$$

$$x = 1$$

$$(1, e^{-1})$$

61. $f'(x) = \ln x + 1 = 0$

$$\ln x = -1$$

$$x = e^{-1}$$

$$(e^{-1}, -e^{-1})$$

62. $f'(x) = \frac{1 - \ln x}{x^2} = 0$

$$1 - \ln x = 0$$

$$x = e$$

$$\left(e, \frac{1}{e}\right)$$


63. (a) $P(t) = -0.63t^2 + 63.3t + 8448$

(c) Using the limit definition,

$$P'(t) = -1.26t + 63.3$$

$$P'(20) = -1.26(20) + 63.3 = 38.1.$$


(d) Answers will vary.




$$P'(20) \approx 38$$

At time 2020, the population is increasing at approximately 38,000 per year.

64. (a) $N = 1.04p^2 - 81.50p + 1613.31$



(d) The rate of decrease in sales decreases as the price increases.



Slope = -50.3 for $p = 15$.

Slope = -19.1 for $p = 30$.

65. (a) $V = \frac{4}{3}\pi r^3$

$$\begin{aligned} V'(r) &= \lim_{h \rightarrow 0} \frac{V(r+h) - V(r)}{h} \\ &= \lim_{h \rightarrow 0} \frac{(4/3)\pi(r+h)^3 - (4/3)\pi r^3}{h} \\ &= \lim_{h \rightarrow 0} \frac{\left(\frac{4}{3}\pi\right) \frac{r^3 + 3r^2h + 3rh^2 + h^3 - r^3}{h}}{h} \\ &= \lim_{h \rightarrow 0} \frac{4}{3}\pi(3r^2 + 3rh + h^2) = 4\pi r^2 \end{aligned}$$

(b) $V'(4) = 4\pi(4)^2 \approx 201.06$

(c) Cubic inches per inch; Answers will vary.

66. $S(r) = 4\pi r^2$

(a) Using the limit definition, $S'(r) = 8\pi r$.

(b) $S'(2) = 8\pi(2) = 16\pi \approx 50.27$

(c) Square millimeters per millimeter

67. $s(t) = -16t^2 + 64t + 80$

(a) Using the limit definition, $s'(t) = -32t + 64$.

(b) $s(0) = 80, s(3) = 128$

$$\begin{aligned} \text{Average rate of change} &= \frac{128 - 80}{3} \\ &= \frac{48}{3} = 16 \text{ ft/sec} \end{aligned}$$


(c) $s'(t) = -32t + 64 = 0 \Rightarrow t = 2$ seconds

Answers will vary.

(d) $s(t) = -16t^2 + 64t + 80$

$$= 0 \Rightarrow t = 5 \text{ seconds}$$

$$s'(5) = -32(5) + 64 = -96 \text{ ft/sec}$$



68. $s(t) = -16t^2 + 120$


(a) Using the limit definition, $s'(t) = -32t$.

$$\begin{aligned} \text{(b) Average} &= \frac{s(2) - s(0)}{2 - 0} \\ &= \frac{56 - 120}{2} = -32 \text{ ft/sec} \end{aligned}$$

(c) $s(t) = -16t^2 + 120 = 0 \Rightarrow t = \sqrt{7.5}$
 $s'(\sqrt{7.5}) = -32\sqrt{7.5} \approx -87.64 \text{ ft/sec}$

(d) $-32t = -60$

$$t = \frac{60}{32} = 1.875 \text{ seconds}$$



69. True. The slope is $2x$, which is different for all x .

70. False. For example, the tangent line to $y = x^3$ at $(1, 1)$ intersects the curve at $(-2, -8)$.


71. Matches (b).
 (Derivative is always positive, but decreasing.)

72. Matches (a).
 (Derivative approaches $-\infty$ when x approaches 0.)


73. Matches (d).
 (Derivative is -1 for $x < 0$, 1 for $x > 0$.)

74. Matches (c).
 (Derivative decreases until origin, then increases.)

75. Answers will vary.



76. Answers not unique




77. $f(x) = \frac{1}{x^2 - x - 2} = \frac{1}{(x + 1)(x - 2)}$

Intercept: $(0, -\frac{1}{2})$

Vertical asymptotes: $x = -1, x = 2$

Horizontal asymptote: $y = 0$




$$78. f(x) = \frac{x-2}{x^2-4x+3} = \frac{x-2}{(x-3)(x-1)}$$

$$\text{Intercepts: } (2, 0), \left(0, -\frac{2}{3}\right)$$

$$\text{Vertical asymptotes: } x = 1, x = 3$$

$$\text{Horizontal asymptote: } y = 0$$




$$79. f(x) = \frac{x^2-x-2}{x-2} = \frac{(x-2)(x+1)}{x-2} = x+1, x \neq 2$$

$$\text{Line with hole at } (2, 3)$$

$$\text{Intercepts: } (0, 1), (-1, 0)$$

$$\text{Slant asymptote: } y = x + 1$$




$$80. f(x) = \frac{x^2-16}{x+4} = \frac{(x+4)(x-4)}{x+4} = x-4, x \neq -4$$

$$\text{Line with hole at } (-4, -8)$$

$$\text{Intercepts: } (0, -4), (4, 0)$$

$$\text{Slant asymptote: } y = x - 4$$



$$81. \langle 1, 1, 1 \rangle \times \langle 2, 1, -1 \rangle = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ 1 & 1 & 1 \\ 2 & 1 & -1 \end{vmatrix} \\ = \langle -2, 3, -1 \rangle$$

$$82. \mathbf{u} \times \mathbf{v} = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ -10 & 0 & 6 \\ 7 & 0 & 0 \end{vmatrix} = \langle 0, 42, 0 \rangle$$

$$83. \langle -4, 10, 0 \rangle \times \langle 4, -1, 0 \rangle = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ -4 & 10 & 0 \\ 4 & -1 & 0 \end{vmatrix} \\ = \langle 0, 0, -36 \rangle$$

$$84. \mathbf{u} \times \mathbf{v} = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ 8 & -7 & 14 \\ -1 & 8 & 4 \end{vmatrix} = \langle -140, -46, 57 \rangle$$

Section 11.4 Limits at Infinity and Limits of Sequences

■ The limit at infinity $\lim_{x \rightarrow \infty} f(x) = L$ means that $f(x)$ get arbitrarily close to L as x increases without bound.

■ Similarly, the limit at infinity $\lim_{x \rightarrow -\infty} f(x) = L$ means that $f(x)$ get arbitrarily close to L as x decreases without bound.

■ You should be able to calculate limits at infinity, especially those arising from rational functions.

■ Limits of functions can be used to evaluate limits of sequences. If f is a function such that $\lim_{x \rightarrow \infty} f(x) = L$ and if a_n is a sequence such that $f(n) = a_n$, then $\lim_{n \rightarrow \infty} a_n = L$.

Vocabulary Check

1. limit, infinity

2. converge

3. diverge

1. Intercept: (0, 0)

Horizontal asymptote: $y = 4$

Matches (c).

2. Horizontal asymptote: $y = 1$

Matches (a).

3. Horizontal asymptote: $y = 4$ Vertical asymptote: $x = 0$

Matches (d).

4. $f(x) = x + \frac{1}{x}$

No horizontal asymptote

Matches (b).

5. Vertical asymptotes: $x = \pm 1$ Horizontal asymptote: $y = 1$

Matches (f).

6. Vertical asymptote: $x = 2$ Horizontal asymptote: $y = 2$


Matches (g).

7. Vertical asymptote: $x = 2$ Horizontal asymptote: $y = -2$


Matches (h).

8. Vertical asymptote: $x = \pm 2$ Horizontal asymptote: $y = -4$


Matches (e).

9. $\lim_{x \rightarrow \infty} \frac{3}{x^2} = 0$ 10. $\lim_{x \rightarrow \infty} \frac{5}{2x} = 0$ 11. $\lim_{x \rightarrow \infty} \frac{3+x}{3-x} = -1$ 12. $\lim_{x \rightarrow \infty} \frac{2-7x}{2+3x} = -\frac{7}{3}$ 13. $\lim_{x \rightarrow -\infty} \frac{5x-2}{6x+1} = \frac{5}{6}$ 14. $\lim_{x \rightarrow -\infty} \frac{5-3x}{x+4} = -3$ 15. $\lim_{x \rightarrow -\infty} \frac{4x^2-3}{2-x^2} = \frac{4}{-1} = -4$ 16. $\lim_{x \rightarrow -\infty} \frac{x^2+3}{5x^2-4} = \frac{1}{5}$ 17. $\lim_{t \rightarrow \infty} \frac{t^2}{t+3}$ does not exist.18. $\lim_{y \rightarrow \infty} \frac{4y^4}{y^2+3}$ does not exist.19. $\lim_{t \rightarrow \infty} \frac{4t^2+3t-1}{3t^2+2t-5} = \frac{4}{3}$ 20. $\lim_{x \rightarrow \infty} \frac{5-6x-3x^2}{2x^2+x+4} = -\frac{3}{2}$ 21. $\lim_{y \rightarrow -\infty} \frac{3+8y-4y^2}{3-y-2y^2} = \frac{-4}{-2} = 2$ 22. $\lim_{t \rightarrow -\infty} \frac{t^2+9t-10}{2+4t-3t^2} = \frac{1}{-3} = -\frac{1}{3}$ 23. $\lim_{x \rightarrow -\infty} \frac{-(x^2+3)}{(2-x)^2} = \lim_{x \rightarrow -\infty} \frac{-x^2-3}{x^2-4x+4} = -1$ 24. $\lim_{x \rightarrow \infty} \frac{2x^2-6}{(x-1)^2} = \lim_{x \rightarrow \infty} \frac{2x^2-6}{x^2-2x+1} = 2$ 25. $\lim_{x \rightarrow -\infty} \left[\frac{x}{(x+1)^2} - 4 \right] = 0 - 4 = -4$ 26. $\lim_{x \rightarrow \infty} \left[7 + \frac{2x^2}{(x+3)^2} \right] = 7 + 2 = 9$ 27. $\lim_{t \rightarrow \infty} \left(\frac{1}{3t^2} - \frac{5t}{t+2} \right) = 0 - 5 = -5$ 28. $\lim_{x \rightarrow \infty} \left[\frac{x}{2x+1} + \frac{3x^2}{(x-3)^2} \right] = \frac{1}{2} + 3 = \frac{7}{2}$ 29. $y = \frac{3x}{1-x}$ Horizontal asymptote: $y = -3$ 


30. $y = \frac{x^2}{x^2 + 4}$

 Horizontal asymptote: $y = 1$



31. $y = \frac{2x}{1 - x^2}$

 Horizontal asymptote: $y = 0$



32. $y = \frac{2x + 1}{x^2 - 1}$

 Horizontal asymptote: $y = 0$


33. $y = 1 - \frac{3}{x^2}$

 Horizontal asymptote: $y = 1$



34. $y = 2 + \frac{1}{x}$

 Horizontal asymptote: $y = 2$


35. $f(x) = x - \sqrt{x^2 + 2}$

(a) x	10^0	10^1	10^2	10^3	10^4	10^5	10^6
$f(x)$	-0.7321	-0.0995	-0.0100	-0.0010	-1.0×10^{-4}	-1.0×10^{-5}	-1.0×10^{-6}

$$\lim_{x \rightarrow \infty} f(x) = 0$$




$$\lim_{x \rightarrow \infty} f(x) = 0$$

36. $f(x) = 3x - \sqrt{9x^2 + 1}$

(a) x	10^0	10^1	10^2	10^3	10^4	10^5	10^6
$f(x)$	-0.1623	-0.0167	-0.0017	-1.67×10^{-4}	-1.67×10^{-5}	-1.67×10^{-6}	-1.67×10^{-7}

$$\lim_{x \rightarrow \infty} f(x) = 0$$




$$\lim_{x \rightarrow \infty} f(x) = 0$$

37. $f(x) = 3(2x - \sqrt{4x^2 + x})$

(a)

x	10^0	10^1	10^2	10^3	10^4	10^5	10^6
$f(x)$	-0.7082	-0.7454	-0.7495	-0.74995	-0.749995	-0.7499995	-0.75

$\lim_{x \rightarrow \infty} f(x) = -0.75$




$\lim_{x \rightarrow \infty} f(x) = -0.75$

38. $f(x) = 4(4x - \sqrt{16x^2 - x})$

(a)

x	10^0	10^1	10^2	10^3	10^4	10^5	10^6
$f(x)$	0.5081	0.5008	0.5001	0.5000	0.5000	0.5000	0.5000

$\lim_{x \rightarrow \infty} f(x) = 0.5$



$\lim_{x \rightarrow \infty} f(x) = 0.5$

39. $a_n = \frac{n+1}{n^2+1}$
 $1, \frac{3}{5}, \frac{2}{5}, \frac{5}{17}, \frac{3}{13}$

$\lim_{n \rightarrow \infty} a_n = 0$

40. $a_n = \frac{n}{n^2+1}$
 $\frac{1}{2}, \frac{2}{5}, \frac{3}{10}, \frac{4}{17}, \frac{5}{26}$

$\lim_{n \rightarrow \infty} \frac{n}{n^2+1} = 0$

41. $a_n = \frac{n}{2n+1}$
 $\frac{1}{3}, \frac{2}{5}, \frac{3}{7}, \frac{4}{9}, \frac{5}{11}$

$\lim_{n \rightarrow \infty} a_n = \frac{1}{2}$

42. $a_n = \frac{4n-1}{n+3}$
 $\frac{3}{4}, \frac{7}{5}, \frac{11}{6}, \frac{15}{7}, \frac{19}{8}$

$\lim_{n \rightarrow \infty} \frac{4n-1}{n+3} = 4$

43. $a_n = \frac{n^2}{3n+2}$
 $\frac{1}{5}, \frac{1}{2}, \frac{9}{11}, \frac{8}{7}, \frac{25}{17}$

$\lim_{n \rightarrow \infty} \frac{n^2}{3n+2}$ does not exist.

44. $a_n = \frac{4n^2+1}{2n}$
 $\frac{5}{2}, \frac{17}{4}, \frac{37}{6}, \frac{65}{8}, \frac{101}{10}$

$\lim_{n \rightarrow \infty} \frac{4n^2+1}{2n}$ does not exist.

45. $a_n = \frac{(n+1)!}{n!}$

2, 3, 4, 5, 6

$\lim_{n \rightarrow \infty} \frac{(n+1)!}{n!} = \lim_{n \rightarrow \infty} (n+1)$ does not exist.

46. $a_n = \frac{(3n-1)!}{(3n+1)!} = \frac{1}{(3n+1)(3n)}$

$\frac{1}{12}, \frac{1}{42}, \frac{1}{90}, \frac{1}{156}, \frac{1}{240}$

$\lim_{n \rightarrow \infty} \frac{(3n-1)!}{(3n+1)!} = 0$

47. $a_n = \frac{(-1)^n}{n}$

$$-1, \frac{1}{2}, -\frac{1}{3}, \frac{1}{4}, -\frac{1}{5}$$

$$\lim_{n \rightarrow \infty} \frac{(-1)^n}{n} = 0$$

48. $a_n = \frac{(-1)^{n+1}}{n^2}$

$$1, -\frac{1}{4}, \frac{1}{9}, -\frac{1}{16}, \frac{1}{25}$$

$$\lim_{n \rightarrow \infty} \frac{(-1)^{n+1}}{n^2} = 0$$

49.

n	10^0	10^1	10^2	10^3	10^4	10^5	10^6
a_n	2	1.55	1.505	1.5005	1.5001	1.500	1.500

$$\lim_{n \rightarrow \infty} a_n = 1.5$$

$$a_n = \frac{1}{n} \left(n + \frac{1}{n} \left[\frac{n(n+1)}{2} \right] \right) = 1 + \frac{1}{n^2} \left[\frac{n^2 + n}{2} \right] = 1 + \frac{n^2 + n}{2n^2}$$

$$\lim_{n \rightarrow \infty} a_n = 1 + \frac{1}{2} = \frac{3}{2}$$

50.

n	10^0	10^1	10^2	10^3	10^4	10^5	10^6
a_n	20	12.8	12.08	12.008	12.0008	12.0001	12.0000

$$\lim_{n \rightarrow \infty} a_n = 12.0$$

$$a_n = \frac{4}{n} \left(n + \frac{4}{n} \left[\frac{n(n+1)}{2} \right] \right) = 4 + \frac{16}{n^2} \left[\frac{n^2 + n}{2} \right] = 4 + \frac{8(n^2 + n)}{n^2}$$

$$\lim_{n \rightarrow \infty} a_n = 4 + 8 = 12$$

51.

n	10^0	10^1	10^2	10^3	10^4	10^5	10^6
a_n	16	6.16	5.4136	5.3413	5.3341	5.3334	5.3333

$$\lim_{n \rightarrow \infty} a_n = 5.33$$

$$a_n = \frac{16}{n^3} \left[\frac{n(n+1)(2n+1)}{6} \right] = \left(\frac{8}{3} \right) \frac{n(n+1)(2n+1)}{n^3}$$

$$\lim_{n \rightarrow \infty} a_n = \frac{16}{3}$$

52.

n	10^0	10^1	10^2	10^3	10^4	10^5	10^6
a_n	1	0.7975	0.7550	0.7505	0.7501	0.7500	0.7500

$$\lim_{n \rightarrow \infty} a_n = 0.75$$

$$a_n = \frac{n(n+1)}{n^2} - \frac{1}{n^4} \left[\frac{n(n+1)}{2} \right]^2 = \frac{n^2 + n}{n^2} - \left(\frac{1}{4} \right) \frac{(n^2 + n)^2}{n^4}$$

$$\lim_{n \rightarrow \infty} a_n = 1 - \frac{1}{4} = \frac{3}{4}$$

53. (a) Average cost = $\bar{C} = \frac{C}{x} = 13.50 + \frac{45,750}{x}$

(b) $\bar{C}(100) = \$471$

$\bar{C}(1000) = \$59.25$

(c) $\lim_{x \rightarrow \infty} \bar{C}(x) = 13.50$

As more units are produced, the fixed costs (45,750) become less dominant.

54. $C = 1.25x + 10,500$


(a) Average cost = $\bar{C} = \frac{C}{x} = 1.25 + \frac{10,500}{x}$

(b) $\bar{C}(100) = \$106.25$

$\bar{C}(1000) = \$11.75$

(c) As $x \rightarrow \infty$, $\bar{C} \rightarrow \$1.25$.

As the number of tons gets very large, the average cost approaches \$1.25 per ton.




(b) For 2004, $t = 14$ and $E(14) \approx 72.0$ million.

For 2008, $t = 18$ and $E(18) \approx 73.8$ million.

(c) $\lim_{t \rightarrow \infty} E(t) = \frac{0.702}{0.009} = 78$

The enrollment approaches 78 million.

(d) Answers will vary.



(b) For 2006, $t = 16$ and $N \approx 2772$ thousand.

For 2010, $t = 20$ and $N \approx 2696$ thousand.

(c) $\lim_{t \rightarrow \infty} N(t) = \frac{40.8189}{0.0157} \approx 2600$

The number of injuries approaches 2600 thousand.

(d) Answers will vary.

57. False. $f(x) = \frac{x^2 + 1}{1}$ does not have a horizontal asymptote.

58. False. The limit does not exist.


59. True

60. False

61. For example, let $f(x) = \frac{1}{x^2}$ and $g(x) = \frac{1}{x^2}$.

Then, $\lim_{x \rightarrow 0} \frac{1}{x^2}$ increases without bound, but

$\lim_{x \rightarrow 0} [f(x) - g(x)] = 0$.







Two horizontal asymptotes: $y = \pm 1$

63. Converges to 0

64. Diverges

65. Diverges

66. Converges to 6




67. $y = x^4$

(a) $f(x) = (x + 3)^4$

(b) $f(x) = x^4 - 1$

(c) $f(x) = -2 + x^4$

(d) $f(x) = \frac{1}{2}(x - 4)^4$




68. $y = x^3$

(a) $f(x) = (x + 2)^3$

(b) $f(x) = 3 + x^3$

(c) $f(x) = 2 - \frac{1}{4}x^3$

(d) $f(x) = 3(x + 1)^3$



69.
$$\begin{array}{r} x^2 + 2x + 1 \\ x^2 - 4 \overline{) x^4 + 2x^3 - 3x^2 - 8x - 4} \\ \underline{x^4 - 4x^2} \\ 2x^3 + x^2 \\ \underline{2x^3 - 8x} \\ x^2 - 4 \end{array}$$

$$2x^3 + x^2$$

$$\underline{2x^3 - 8x}$$

$$x^2 - 4$$

$$x^4 + 2x^3 - 3x^2 - 8x - 4 = (x^2 - 4)(x^2 + 2x + 1)$$

70.
$$\begin{array}{r} 2x^3 + 4x^2 - 2x - 8 \\ x^2 - 2x + 1 \overline{) 2x^5 - 8x^3 + 4x - 1} \\ \underline{2x^5 - 4x^4 + 2x^3} \\ 4x^4 - 10x^3 \\ \underline{4x^4 - 8x^3 + 4x^2} \\ -2x^3 - 4x^2 + 4x \\ \underline{-2x^3 + 4x^2 - 2x} \\ -8x^2 + 6x - 1 \\ \underline{-8x^2 + 16x - 8} \\ -10x + 7 \end{array}$$

$$2x^5 - 4x^4 + 2x^3$$

$$4x^4 - 10x^3$$

$$\underline{4x^4 - 8x^3 + 4x^2}$$

$$-2x^3 - 4x^2 + 4x$$

$$\underline{-2x^3 + 4x^2 - 2x}$$

$$-8x^2 + 6x - 1$$

$$\underline{-8x^2 + 16x - 8}$$

$$-10x + 7$$

$$\frac{2x^5 - 8x^3 + 4x - 1}{x^2 - 2x + 1} = 2x^3 + 4x^2 - 2x - 8 + \frac{-10x + 7}{x^2 - 2x + 1}$$

71.
$$\begin{array}{r} x^3 + 5x^2 - 3 \\ 3x + 2 \overline{) 3x^4 + 17x^3 + 10x^2 - 9x - 8} \\ \underline{3x^4 + 2x^3} \\ 15x^3 + 10x^2 \\ \underline{15x^3 + 10x^2} \\ -9x - 8 \\ \underline{-9x - 6} \\ -2 \end{array}$$

$$3x^4 + 2x^3$$

$$15x^3 + 10x^2$$

$$\underline{15x^3 + 10x^2}$$

$$-9x - 8$$

$$\underline{-9x - 6}$$

$$-2$$

$$\frac{3x^4 + 17x^3 + 10x^2 - 9x - 8}{3x + 2} = x^3 + 5x^2 - 3 + \frac{-2}{3x + 2}$$

$$72. \frac{2x^2 + 11x + 14}{5x - 2} \div \frac{10x^3 + 51x^2 + 48x - 28}{10x^3 - 4x^2}$$

$$= \frac{55x^2 + 48x}{55x^2 - 22x} \cdot \frac{70x - 28}{70x - 28}$$


$$\frac{10x^3 + 51x^2 + 48x - 28}{5x - 2} = 2x^2 + 11x + 14, \quad x \neq \frac{2}{5}$$

$$73. f(x) = x^4 - x^3 - 20x^2$$

$$= x^2(x^2 - x - 20)$$

$$= x^2(x - 5)(x + 4)$$


Real zeros: 0, 0, 5, -4



$$74. x^5 + x^3 - 6x = x(x^4 + x^2 - 6)$$

$$= x(x^2 + 3)(x^2 - 2)$$

Real zeros: 0, $\pm\sqrt{2}$




$$75. f(x) = x^3 - 3x^2 + 2x - 6$$

$$= x^2(x - 3) + 2(x - 3)$$

$$= (x - 3)(x^2 + 2)$$


Real zero: 3



$$76. x^3 - 4x^2 - 25x + 100 = x^2(x - 4) - 25(x - 4)$$

$$= (x^2 - 25)(x - 4)$$

Real zeros: $\pm 5, 4$



$$77. \sum_{i=1}^6 (2i + 3) = 5 + 7 + 9 + 11 + 13 + 15 = 60$$

$$78. \sum_{i=0}^4 5i^2 = 0 + 5 + 20 + 45 + 80 = 150$$

$$79. \sum_{k=1}^{10} 15 = 10(15) = 150$$

$$80. \sum_{k=0}^8 \frac{3}{k^2 + 1} \approx 5.8791$$

Section 11.5 The Area Problem

■ You should know the following summation formulas and properties.

$$(a) \sum_{i=1}^n c = cn$$

$$(b) \sum_{i=1}^n i = \frac{n(n+1)}{2}$$

$$(c) \sum_{i=1}^n i^2 = \frac{n(n+1)(2n+1)}{6}$$

$$(d) \sum_{i=1}^n i^3 = \frac{n^2(n+1)^2}{4}$$

$$(e) \sum_{i=1}^n (a_i \pm b_i) = \sum_{i=1}^n a_i \pm \sum_{i=1}^n b_i$$

$$(f) \sum_{i=1}^n ka_i = k \sum_{i=1}^n a_i$$

■ You should be able to evaluate a limit of a summation, $\lim_{n \rightarrow \infty} S(n)$.

■ You should be able to approximate the area of a region using rectangles. By increasing the number of rectangles, the approximation improves.

—CONTINUED—

—CONTINUED—

- The area of a plane region above the x -axis bounded by f between $x = a$ and $x = b$ is the limit of the sum of the approximating rectangles:

$$A = \lim_{n \rightarrow \infty} \sum_{i=1}^n f\left(a + \frac{(b-a)i}{n}\right) \left(\frac{b-a}{n}\right)$$

- You should be able to use the limit definition of area to find the area bounded by simple functions in the plane.

Vocabulary Check

1. $\frac{n(n+1)}{2}$

2. $\frac{n^2(n+1)^2}{4}$

3. area

1. $\sum_{i=1}^{60} 7 = 7(60) = 420$

2. $\sum_{i=1}^{45} 3 = 3(45) = 135$

3. $\sum_{i=1}^{20} i^3 = \frac{20^2(21)^2}{4} = 44,100$

4. $\sum_{i=1}^{30} i^2 = \frac{n(n+1)(2n+1)}{6}$
 $= \frac{30(31)(61)}{6} = 9455$

5. $\sum_{k=1}^{20} (k^3 + 2) = \frac{20^2(21)^2}{4} + 2(20)$
 $= 44,100 + 40$
 $= 44,140$

6. $\sum_{k=1}^{50} (2k+1) = 2 \sum_{k=1}^{50} k + \sum_{k=1}^{50} 1$
 $= 2 \frac{50(51)}{2} + 50$
 $= 2600$

7. $\sum_{j=1}^{25} (j^2 + j) = \frac{25(26)(51)}{6} + \frac{25(26)}{2} = 5850$

8. $\sum_{j=1}^{10} (j^3 - 3j^2) = \frac{10^2(11)^2}{4} - 3 \left(\frac{10(11)(21)}{6} \right) = 1870$

9. (a) $S(n) = \sum_{i=1}^n \frac{i^3}{n^4} = \frac{1}{n^4} \left[\frac{n^2(n+1)^2}{4} \right] = \frac{n^2 + 2n + 1}{4n^2}$

(b)	n	10^0	10^1	10^2	10^3	10^4
	$S(n)$	1	0.3025	0.255025	0.25050025	0.25005

(c) $\lim_{n \rightarrow \infty} S(n) = \frac{1}{4}$

10. (a) $S(n) = \sum_{i=1}^n \frac{i}{n^2} = \frac{1}{n^2} \frac{n(n+1)}{2} = \frac{n+1}{2n}$

(b)	n	10^0	10^1	10^2	10^3	10^4
	$S(n)$	1	0.55	0.505	0.5005	0.50005

(c) $\lim_{n \rightarrow \infty} S(n) = \frac{1}{2}$

11. (a) $S(n) = \sum_{i=1}^n \frac{3}{n^3} (1+i^2) = \frac{3}{n^3} \left[n + \frac{n(n+1)(2n+1)}{6} \right] = \frac{3}{n^2} + \frac{6n^2 + 9n + 3}{6n^2} = \frac{2n^2 + 3n + 7}{2n^2}$

(b)	n	10^0	10^1	10^2	10^3	10^4
	$S(n)$	6	1.185	1.0154	1.0015	1.00015

(c) $\lim_{n \rightarrow \infty} S(n) = 1$

12. (a) $S(n) = \sum_{i=1}^n \frac{2i + 3}{n^2} = \frac{1}{n^2} \left(2 \left(\frac{n(n+1)}{2} \right) + 3n \right) = \frac{n+1}{n} + \frac{3}{n} = \frac{n+4}{n}$

(b)

n	10^0	10^1	10^2	10^3	10^4
$S(n)$	5	1.4	1.04	1.004	1.0004

(c) $\lim_{n \rightarrow \infty} S(n) = 1$

13. (a) $S(n) = \sum_{i=1}^n \left(\frac{i^2}{n^3} + \frac{2}{n} \right) \left(\frac{1}{n} \right) = \frac{1}{n} \left[\frac{n(n+1)(2n+1)}{6n^3} + \frac{2n}{n} \right] = \frac{1}{6n^3} (2n^2 + 3n + 1) + \frac{2}{n} = \frac{14n^2 + 3n + 1}{6n^3}$

(b)

n	10^0	10^1	10^2	10^3	10^4
$S(n)$	3	0.2385	0.02338	0.00233	0.0002333

(c) $\lim_{n \rightarrow \infty} S(n) = 0$

14. (a) $S(n) = \sum_{i=1}^n \left[3 - 2 \left(\frac{i}{n} \right) \right] \frac{1}{n} = \frac{1}{n} \left[3n - \frac{2n(n+1)}{2} \right] = 3 - \frac{n+1}{n} = \frac{2n-1}{n}$

(b)

n	10^0	10^1	10^2	10^3	10^4
$S(n)$	1	1.9	1.99	1.999	1.9999

(c) $\lim_{n \rightarrow \infty} S(n) = 2$

15. (a) $S(n) = \sum_{i=1}^n \left[1 - \left(\frac{i}{n} \right)^2 \right] \left(\frac{1}{n} \right) = \frac{1}{n} \left[n - \frac{1}{n^2} \left(\frac{n(n+1)(2n+1)}{6} \right) \right] = 1 - \frac{2n^2 + 3n + 1}{6n^2} = \frac{4n^2 - 3n - 1}{6n^2}$

(b)

n	10^0	10^1	10^2	10^3	10^4
$S(n)$	0	0.615	0.66165	0.66617	0.666617

(c) $\lim_{n \rightarrow \infty} S(n) = \frac{2}{3}$

16. (a) $S(n) = \sum_{i=1}^n \left(\frac{4}{n} + \frac{2i}{n^2} \right) \left(\frac{2i}{n} \right) = \frac{2}{n} \left[\frac{4n(n+1)}{2} + \frac{2n(n+1)(2n+1)}{6} \right]$
 $= \frac{2}{n} \left[\frac{4n^2 + 4n}{2} + \frac{2(2n^2 + 3n + 1)}{6} \right] = \frac{16n^2 + 18n + 2}{3n^2}$

(b)

n	10^0	10^1	10^2	10^3	10^4
$S(n)$	12.0	5.94	5.3934	5.3393	5.33393

(c) $\lim_{n \rightarrow \infty} S(n) = \frac{16}{3}$

17. $f(x) = x + 4, [-1, 2], n = 6, \text{width} = \frac{1}{2}$
 Area $\approx \frac{1}{2} [3.5 + 4 + 4.5 + 5 + 5.5 + 6] = 14.25$ square units

18. $f(x) = 2 - x^2, -1 \leq x \leq 1, n = 4, \text{width} = \frac{1}{2}$
 Area $\approx \frac{1}{2} \left[\left(2 - \left(-\frac{1}{2} \right)^2 \right) + (2 - 0^2) + \left(2 - \left(\frac{1}{2} \right)^2 \right) + (2 - 1^2) \right]$
 $= \frac{1}{2} [1.75 + 2 + 1.75 + 1]$
 $= 3.25$ square units

19. The width of each rectangle is $\frac{1}{4}$. The height is obtained by evaluating f at the right-hand endpoint of each interval.

$$A \approx \sum_{i=1}^8 f\left(\frac{i}{4}\right)\left(\frac{1}{4}\right) = \sum_{i=1}^8 \frac{1}{4}\left(\frac{i}{4}\right)^3\left(\frac{1}{4}\right) = 1.265625 \text{ square units}$$

20. Area $\approx \frac{1}{2}\left[\frac{1}{16} + \frac{1}{2} + \frac{27}{16} + 4\right] = 3.125$ square units

21. Width of each rectangle is $12/n$. The height is

$$f\left(\frac{12i}{n}\right) = -\frac{1}{3}\left(\frac{12i}{n}\right) + 4.$$

$$A \approx \sum_{i=1}^n \left[-\frac{1}{3}\left(\frac{12i}{n}\right) + 4\right]\left(\frac{12}{n}\right)$$

Note: Exact area is 24.

n	4	8	20	50
Approximate area	18	21	22.8	23.52

22. The width of each rectangle is $3/n$. The height is

$$f\left(\frac{3i}{n}\right) = 9 - \left(\frac{3i}{n}\right)^2.$$

$$A \approx \sum_{i=1}^n \left(9 - \left(\frac{3i}{n}\right)^2\right)\frac{3}{n}$$

Note: Exact area is 18.

n	4	8	20	50
Approximate area	14.344	16.242	17.314	17.7282

23. The width of each rectangle is $3/n$. The height is

$$\frac{1}{9}\left(\frac{3i}{n}\right)^3.$$

$$A \approx \sum_{i=1}^n \frac{1}{9}\left(\frac{3i}{n}\right)^3\left(\frac{3}{n}\right)$$

n	4	8	20	50
Approximate area	3.52	2.85	2.48	2.34

24. The width of each rectangle is $(2 - (-1))/n = 3/n$. The height is

$$f\left(-1 + \frac{3i}{n}\right) = 3 - \frac{1}{4}\left(-1 + \frac{3i}{n}\right)^3.$$

$$A \approx \sum_{i=1}^n \left[3 - \frac{1}{4}\left(-1 + \frac{3i}{n}\right)^3\right]\frac{3}{n}$$

Note: Exact area is $8\frac{1}{16} = 8.0625$.

n	4	8	20	50
Approximate area	7.113	7.614	7.8895	7.994

25. $f(x) = 2x + 5$, $[0, 4]$

The width of each rectangle is $4/n$. The height is

$$f\left(\frac{4i}{n}\right) = 2\left(\frac{4i}{n}\right) + 5 = \frac{8i}{n} + 5.$$

$$A \approx \sum_{i=1}^n \left(\frac{8i}{n} + 5\right)\left(\frac{4}{n}\right)$$

n	4	8	20	50	100	∞
Area	40	38	36.8	36.32	36.16	36

25. —CONTINUED—

$$A \approx \sum_{i=1}^n \left(\frac{8i}{n} + 5 \right) \left(\frac{4}{n} \right) = \sum_{i=1}^n \left(\frac{20}{n} + \frac{32}{n^2} i \right) \approx \frac{20}{n}(n) + \frac{32}{n^2} \left(\frac{n(n+1)}{2} \right) = 20 + 16 \left(\frac{n^2+n}{n^2} \right)$$

$$A = \lim_{n \rightarrow \infty} \left[20 + 16 \left(\frac{n^2+n}{n^2} \right) \right] = 20 + 16 = 36$$

26. $f(x) = 3x + 1$, $[0, 4]$

The width of each rectangle is $4/n$. The height is

$$f\left(\frac{4i}{n}\right) = 3\left(\frac{4i}{n}\right) + 1.$$

n	4	8	20	50	100	∞
Area	34	31	29.2	28.48	28.24	28

$$A \approx \sum_{i=1}^n \left(\frac{12i}{n} + 1 \right) \frac{4}{n}$$

$$A \approx \sum_{i=1}^n \left(\frac{48i}{n^2} + \frac{4}{n} \right) = \frac{48}{n^2} \left(\frac{n(n+1)}{2} \right) + \frac{4}{n}(n) = 24 \left(\frac{n^2+n}{n^2} \right) + 4$$

$$A = \lim_{n \rightarrow \infty} \left[24 \left(\frac{n^2+n}{n^2} \right) + 4 \right] = 28$$

27. $f(x) = 16 - 2x$, $[1, 5]$

The width of each rectangle is $4/n$. The height is

$$f\left(1 + \frac{4i}{n}\right) = 16 - 2\left(1 + \frac{4i}{n}\right) = 14 - \frac{8i}{n}.$$

n	4	8	20	50	100	∞
Area	36	38	39.2	39.68	39.84	40

$$A \approx \sum_{i=1}^n \left(14 - \frac{8i}{n} \right) \frac{4}{n}$$

$$A \approx \sum_{i=1}^n \left(\frac{56}{n} - \frac{32i}{n^2} \right) = \frac{56}{n}(n) - \frac{32}{n^2} \left(\frac{n(n+1)}{2} \right) = 56 - 16 \left(\frac{n(n+1)}{n^2} \right)$$

$$A = \lim_{n \rightarrow \infty} \left[56 - 16 \left(\frac{n(n+1)}{n^2} \right) \right] = 56 - 16 = 40$$

28. $f(x) = 20 - 2x$, $[2, 6]$

The width of each rectangle is $4/n$. The height is

$$f\left(2 + \frac{4i}{n}\right) = 20 - 2\left(2 + \frac{4i}{n}\right) = 16 - \frac{8i}{n}.$$

n	4	8	20	50	100	∞
Area	44	46	47.2	47.68	47.84	48

$$A \approx \sum_{i=1}^n \left(16 - \frac{8i}{n} \right) \frac{4}{n}$$

$$A \approx \sum_{i=1}^n \left(\frac{64}{n} - \frac{32i}{n^2} \right) = \frac{64}{n}(n) - \frac{32}{n^2} \left(\frac{n(n+1)}{2} \right) = 64 - 16 \left(\frac{n(n+1)}{n^2} \right)$$

$$A = \lim_{n \rightarrow \infty} \left[64 - 16 \left(\frac{n(n+1)}{n^2} \right) \right] = 64 - 16 = 48$$

29. $f(x) = 9 - x^2$, $[0, 2]$

The width of each rectangle is $2/n$. The height is

$$f\left(\frac{2i}{n}\right) = 9 - \left(\frac{2i}{n}\right)^2 = 9 - \frac{4i^2}{n^2}.$$

$$A \approx \sum_{i=1}^n \left(9 - \frac{4i^2}{n^2}\right) \left(\frac{2}{n}\right)$$

n	4	8	20	50	100	∞
Area	14.25	14.8125	15.13	15.2528	15.2932	$\frac{46}{3}$

$$A \approx \sum_{i=1}^n \left(\frac{18}{n} - \frac{8i^2}{n^3}\right) = \left(\frac{18}{n}\right)n - \frac{8}{n^3} \left[\frac{n(n+1)(2n+1)}{6}\right] = 18 - \frac{4}{3} \left[\frac{n(n+1)(2n+1)}{n^3}\right]$$

$$A = \lim_{n \rightarrow \infty} \left[18 - \frac{4}{3} \left(\frac{n(n+1)(2n+1)}{n^3}\right)\right] = 18 - \frac{8}{3} = \frac{46}{3}$$

30. $f(x) = x^2 + 1$, $[4, 6]$

The width of each rectangle is $2/n$. The height is

$$f\left(4 + \frac{2i}{n}\right) = \left(4 + \frac{2i}{n}\right)^2 + 1 = 17 + \frac{16i}{n} + \frac{4i^2}{n^2}.$$

$$A \approx \sum_{i=1}^n \left(17 + \frac{16i}{n} + \frac{4i^2}{n^2}\right) \left(\frac{2}{n}\right)$$

n	4	8	20	50	100	∞
Area	57.75	55.1875	53.67	53.0672	52.8668	$\frac{158}{3}$

$$A \approx \sum_{i=1}^n \left(\frac{34}{n} + \frac{32i}{n^2} + \frac{8i^2}{n^3}\right) = \left(\frac{34}{n}\right)n + \left(\frac{32}{n^2}\right)\frac{n(n+1)}{2} + \left(\frac{8}{n^3}\right)\frac{n(n+1)(2n+1)}{6}$$

$$= 34 + 16\left(\frac{n^2+n}{n^2}\right) + \frac{4}{3} \left[\frac{n(n+1)(2n+1)}{n^3}\right]$$

$$A = \lim_{n \rightarrow \infty} \left[34 + 16\left(\frac{n^2+n}{n^2}\right) + \frac{4}{3} \left(\frac{n(n+1)(2n+1)}{n^3}\right)\right] = 34 + 16 + \frac{8}{3} = \frac{158}{3}$$

31. $f(x) = \frac{1}{2}x + 4$, $[-1, 3]$

The width of each rectangle is $4/n$. The height is

$$f\left(-1 + \frac{4i}{n}\right) = \frac{1}{2}\left(-1 + \frac{4i}{n}\right) + 4 = \frac{7}{2} + \frac{2i}{n}.$$

$$A \approx \sum_{i=1}^n \left(\frac{7}{2} + \frac{2i}{n}\right) \left(\frac{4}{n}\right)$$

$$A \approx \sum_{i=1}^n \left(\frac{14}{n} + \frac{8i}{n^2}\right) = \left(\frac{14}{n}\right)n + \frac{8}{n^2} \left(\frac{n(n+1)}{2}\right)$$

$$A = \lim_{n \rightarrow \infty} \left[14 + \frac{4}{n^2} \left(\frac{n(n+1)}{1}\right)\right] = 14 + 4 = 18$$

n	4	8	20	50	100	∞
Area	19	18.5	18.2	18.08	18.04	18

$$32. f(x) = \frac{1}{2}x + 1, [-2, 2]$$

The width of each rectangle is $4/n$. The height is

$$f\left(-2 + \frac{4i}{n}\right) = \frac{1}{2}\left(-2 + \frac{4i}{n}\right) + 1 = \frac{2i}{n}.$$

$$A \approx \sum_{i=1}^n \left(\frac{2i}{n}\right)\left(\frac{4}{n}\right) = \sum_{i=1}^n \frac{8i}{n^2}$$

$$A \approx \sum_{i=1}^n \frac{8i}{n^2} = \frac{8}{n^2} \left(\frac{n(n+1)}{2}\right)$$

$$A = \lim_{n \rightarrow \infty} \left[\frac{8}{n^2} \left(\frac{n(n+1)}{2}\right) \right] = 4$$

n	4	8	20	50	100	∞
Area	5	4.5	4.2	4.08	4.04	4

$$\begin{aligned} 33. A &\approx \sum_{i=1}^n f\left(\frac{i}{n}\right)\left(\frac{1}{n}\right) \\ &= \sum_{i=1}^n \left[4\left(\frac{i}{n}\right) + 1\right]\left(\frac{1}{n}\right) \\ &= \frac{1}{n} \sum_{i=1}^n \left[\frac{4}{n}i + 1\right] \\ &= \frac{1}{n} \left[\frac{4}{n} \frac{n(n+1)}{2} + n \right] \\ &= \frac{1}{n} [2(n+1) + n] \\ &= \frac{3n+2}{n} \end{aligned}$$

$$A = \lim_{n \rightarrow \infty} \frac{3n+2}{n} = 3 \text{ square units}$$

$$\begin{aligned} 35. A &\approx \sum_{i=1}^n f\left(\frac{i}{n}\right)\left(\frac{1}{n}\right) \\ &= \sum_{i=1}^n \left[-2\left(\frac{i}{n}\right) + 3\right]\left(\frac{1}{n}\right) \\ &= \frac{1}{n} \sum_{i=1}^n \left[-\frac{2i}{n} + 3\right] \\ &= \frac{1}{n} \left[-\frac{2}{n} \frac{n(n+1)}{2} + 3n \right] \\ &= \frac{1}{n} [2n - 1] \end{aligned}$$

$$A = \lim_{n \rightarrow \infty} \frac{2n-1}{n} = 2 \text{ square units}$$

$$\begin{aligned} 34. A &\approx \sum_{i=1}^n f\left(\frac{2i}{n}\right)\left(\frac{2}{n}\right) \\ &= \sum_{i=1}^n \left[3\left(\frac{2i}{n}\right) + 2\right]\frac{2}{n} \\ &= \frac{2}{n} \sum_{i=1}^n \left(\frac{6}{n}i + 2\right) \\ &= \frac{2}{n} \left[\frac{6}{n} \frac{n(n+1)}{2} + 2n \right] \\ &= 6\left(\frac{n+1}{n}\right) + 4 \end{aligned}$$

$$A = \lim_{n \rightarrow \infty} \left[6\frac{n+1}{n} + 4 \right] = 10 \text{ square units}$$

$$\begin{aligned} 36. A &\approx \sum_{i=1}^n f\left(2 + \frac{3i}{n}\right)\left(\frac{3}{n}\right) \\ &= \sum_{i=1}^n \left[3\left(2 + \frac{3i}{n}\right) - 4\right]\frac{3}{n} \\ &= \frac{3}{n} \sum_{i=1}^n \left[2 + \frac{9}{n}i\right] \\ &= \frac{3}{n} \left[2n + \frac{9}{n} \frac{n(n+1)}{2} \right] \\ &= 6 + \frac{27}{2} \frac{(n+1)}{n} \end{aligned}$$

$$A = \lim_{n \rightarrow \infty} \left[6 + \frac{27}{2} \frac{(n+1)}{n} \right] = \frac{39}{2} \text{ square units}$$

$$\begin{aligned}
37. A &\approx \sum_{i=1}^n f\left(-1 + \frac{2i}{n}\right)\left(\frac{2}{n}\right) \\
&= \sum_{i=1}^n \left[2 - \left(-1 + \frac{2i}{n}\right)^2\right] \frac{2}{n} \\
&= \sum_{i=1}^n \left[2 - 1 + \frac{4i}{n} - \frac{4i^2}{n^2}\right] \left(\frac{2}{n}\right) \\
&= \frac{2}{n} \sum_{i=1}^n 1 + \frac{8}{n^2} \sum_{i=1}^n i - \frac{8}{n^3} \sum_{i=1}^n i^2 \\
&= \frac{2}{n}(n) + \frac{8}{n^2} \frac{n(n+1)}{2} - \frac{8}{n^3} \frac{n(n+1)(2n+1)}{6} \\
A &= \lim_{n \rightarrow \infty} \left[2 + 4 \frac{n(n+1)}{n^2} - \frac{4}{3} \frac{n(n+1)(2n+1)}{n^3}\right] \\
&= 2 + 4 - \frac{8}{3} = \frac{10}{3} \text{ square units}
\end{aligned}$$

$$\begin{aligned}
38. A &\approx \sum_{n=1}^{\infty} f\left(\frac{i}{n}\right)\left(\frac{1}{n}\right) \\
&= \sum_{n=1}^{\infty} \left[\left(\frac{i}{n}\right)^2 + 2\right] \frac{1}{n} \\
&= \frac{1}{n} \left[\frac{1}{n^2} \frac{n(n+1)(2n+1)}{6} + 2n\right] \\
&= \frac{(n+1)(2n+1)}{6n^2} + 2 \\
A &= \lim_{n \rightarrow \infty} \left[\frac{(n+1)(2n+1)}{6n^2} + 2\right] = \frac{7}{3} \text{ square units}
\end{aligned}$$

$$\begin{aligned}
39. A &\approx \sum_{i=1}^n g\left(1 + \frac{i}{n}\right)\left(\frac{1}{n}\right) \\
&= \sum_{i=1}^n \left[8 - \left(1 + \frac{i}{n}\right)^3\right] \frac{1}{n} \\
&= \sum_{i=1}^n \left[7 - \frac{3i}{n} - \frac{3i^2}{n^2} - \frac{i^3}{n^3}\right] \frac{1}{n} \\
&= \frac{7}{n} \sum_{i=1}^n 1 - \frac{3}{n^2} \sum_{i=1}^n i - \frac{3}{n^3} \sum_{i=1}^n i^2 - \frac{1}{n^4} \sum_{i=1}^n i^3 \\
&= \frac{7}{n}(n) - \frac{3}{n^2} \frac{n(n+1)}{2} - \frac{3}{n^3} \frac{n(n+1)(2n+1)}{6} - \frac{1}{n^4} \frac{n^2(n+1)^2}{4} \\
A &= \lim_{n \rightarrow \infty} \left[7 - \frac{3}{2} \frac{n(n+1)}{n^2} - \frac{1}{2n^3} n(n+1)(2n+1) - \frac{1}{n^4} \frac{n^2(n+1)^2}{4}\right] = 7 - \frac{3}{2} - 1 - \frac{1}{4} = \frac{17}{4} \text{ square units}
\end{aligned}$$

$$\begin{aligned}
40. A &\approx \sum_{i=1}^n g\left(1 + \frac{3i}{n}\right)\left(\frac{3}{n}\right) \\
&= \sum_{i=1}^n \left[64 - \left(1 + \frac{3i}{n}\right)^3\right] \frac{3}{n} \\
&= \frac{3}{n} \sum_{i=1}^n \left[63 - \frac{9i}{n} - \frac{27i^2}{n^2} - \frac{27i^3}{n^3}\right] \\
&= \frac{3}{n} \left[63n - \frac{9n(n+1)}{2n} - \frac{27n(n+1)(2n+1)}{6n^2} - \frac{27n^2(n+1)^2}{4n^3}\right] \\
&= 189 - \frac{27(n+1)}{2n} - \frac{27(n+1)(2n+1)}{2n^2} - \frac{81(n+1)^2}{4n^2} \\
A &= \lim_{n \rightarrow \infty} \left[189 - \frac{27(n+1)}{2n} - \frac{27(n+1)(2n+1)}{2n^2} - \frac{81(n+1)^2}{4n^2}\right] = \frac{513}{4} \text{ square units}
\end{aligned}$$

$$\begin{aligned}
41. A &\approx \sum_{i=1}^n g\left(\frac{i}{n}\right)\left(\frac{1}{n}\right) \\
&= \sum_{i=1}^n \left[2\left(\frac{i}{n}\right) - \left(\frac{i}{n}\right)^3\right]\left(\frac{1}{n}\right) \\
&= \frac{1}{n} \sum_{i=1}^n \left[\frac{2i}{n} - \frac{1}{n^3}i^3\right] \\
&= \frac{1}{n} \left[\frac{2n(n+1)}{2} - \frac{1}{n^3} \frac{n^2(n+1)^2}{4} \right] \\
&= \frac{n+1}{n} - \frac{(n+1)^2}{4n^2} \\
A &= \lim_{n \rightarrow \infty} \left[\frac{n+1}{n} - \frac{(n+1)^2}{4n^2} \right] \\
&= 1 - \frac{1}{4} = \frac{3}{4} \text{ square units}
\end{aligned}$$

$$\begin{aligned}
42. A &\approx \sum_{i=1}^n g\left(\frac{2i}{n}\right)\left(\frac{2}{n}\right) \\
&= \sum_{i=1}^n \left[4\left(\frac{2i}{n}\right) - \left(\frac{2i}{n}\right)^3\right]\left(\frac{2}{n}\right) \\
&= \frac{16}{n^2} \sum_{i=1}^n i - \frac{16}{n^4} \sum_{i=1}^n i^3 \\
&= \frac{16n(n+1)}{n^2} - \frac{16n^2(n+1)^2}{n^4} \\
A &= \lim_{n \rightarrow \infty} \left[8 \frac{n(n+1)}{n^2} - 4 \frac{n^2(n+1)^2}{n^4} \right] \\
&= 8 - 4 = 4 \text{ square units}
\end{aligned}$$


$$\begin{aligned}
43. A &\approx \sum_{i=1}^n f\left(1 + \frac{3i}{n}\right)\left(\frac{3}{n}\right) \\
&= \sum_{i=1}^n \left[\frac{1}{4} \left(1 + \frac{3i}{n}\right)^2 + \left(1 + \frac{3i}{n}\right) \right] \left(\frac{3}{n}\right) \\
&= \sum_{i=1}^n \left(\frac{1}{4} + \frac{3i}{2n} + \frac{9i^2}{4n^2} + 1 + \frac{3i}{n} \right) \left(\frac{3}{n}\right) \\
&= \frac{15}{4n} \sum_{i=1}^n 1 + \frac{27}{2n^2} \sum_{i=1}^n i + \frac{27}{4n^3} \sum_{i=1}^n i^2 \\
&= \frac{15}{4n}(n) + \frac{27}{2n^2} \left(\frac{n(n+1)}{2} \right) + \frac{27}{4n^3} \frac{n(n+1)(2n+1)}{6} \\
A &= \lim_{n \rightarrow \infty} \left[\frac{15}{4} + \frac{27}{4} \frac{n(n+1)}{n^2} + \frac{9}{8n^3} n(n+1)(2n+1) \right] \\
&= \frac{15}{4} + \frac{27}{4} + \frac{9}{4} = \frac{51}{4} \text{ square units}
\end{aligned}$$

$$\begin{aligned}
44. A &\approx \sum_{i=1}^n f\left(-1 + \frac{2i}{n}\right)\left(\frac{2}{n}\right) \\
&= \sum_{i=1}^n \left[\left(-1 + \frac{2i}{n}\right)^2 - \left(-1 + \frac{2i}{n}\right)^3 \right] \left(\frac{2}{n}\right) \\
&= \sum_{i=1}^n \left[\left(1 - \frac{4i}{n} + \frac{4i^2}{n^2}\right) - \left(-1 + \frac{6i}{n} - \frac{12i^2}{n^2} + \frac{8i^3}{n^3}\right) \right] \frac{2}{n} \\
&= \sum_{i=1}^n \left[2 - \frac{10i}{n} + \frac{16i^2}{n^2} - \frac{8i^3}{n^3} \right] \frac{2}{n} \\
&= \frac{4}{n} \sum_{i=1}^n 1 - \frac{20}{n^2} \sum_{i=1}^n i + \frac{32}{n^3} \sum_{i=1}^n i^2 - \frac{16}{n^4} \sum_{i=1}^n i^3 \\
&= \frac{4}{n}(n) - \frac{20}{n^2} \frac{n(n+1)}{2} + \frac{32}{n^3} \frac{n(n+1)(2n+1)}{6} - \frac{16}{n^4} \frac{n^2(n+1)^2}{4} \\
\lim_{n \rightarrow \infty} A &= 4 - 10 + \frac{32}{3} - 4 = \frac{2}{3} \text{ square units}
\end{aligned}$$


45. $y = (-3.0 \cdot 10^{-6})x^3 + 0.002x^2 - 1.05x + 400$

Note that $y = 0$ when $x = 500$.

Area $\approx 105,208.33$ square feet ≈ 2.4153 acres



46. (a) $-4.089 \times 10^{-5}x^3 + 0.01615x^2 - 2.6716x + 452.9286$



(c) Using a graphing utility to integrate from 0 to 300 gives Area $\approx 78,204$ square feet.

(Answers will vary.)

47. True. See Formula 2, page 820.

48. False. n approaches infinity.

49. Answers will vary.

50. Area is approximately a triangle of base 2 and height 3.

Area ≈ 4 ; (c)

51. $2 \tan x = \tan 2x = \frac{2 \tan x}{1 - \tan^2 x}$

$$\tan x = 0 \Rightarrow x = n\pi$$

52. $\cos 2x - 3 \sin x = 2$

$$1 - 2 \sin^2 x - 3 \sin x = 2$$

$$2 \sin^2 x + 3 \sin x + 1 = 0$$

$$(2 \sin x + 1)(\sin x + 1) = 0$$

$$\sin x = \frac{-1}{2} \Rightarrow x = \frac{7\pi}{6} + 2n\pi, \frac{11\pi}{6} + 2n\pi$$

$$\sin x = -1 \Rightarrow x = \frac{3\pi}{2} + 2n\pi$$

53. $(\mathbf{u} \cdot \mathbf{v})\mathbf{u} = (\langle 4, -5 \rangle \cdot \langle -1, -2 \rangle)\langle 4, -5 \rangle$
 $= 6\langle 4, -5 \rangle$
 $= \langle 24, -30 \rangle$

54. $3\mathbf{u} \cdot \mathbf{v} = 3\langle 4, -5 \rangle \cdot \langle -1, -2 \rangle = 3(-4 + 10)$
 $= 18$

55. $\|\mathbf{v}\| - 2 = \sqrt{5} - 2$

56. $\|\mathbf{u}\|^2 - \|\mathbf{v}\|^2 = (4^2 + (-5)^2) - ((-1)^2 + (-2)^2)$
 $= (16 + 25) - (1 + 4)$
 $= 36$

Review Exercises for Chapter 11

1. $\lim_{x \rightarrow 3} (6x - 1)$

The limit (17) can be reached.

x	2.9	2.99	2.999	3	3.001	3.01	3.1
$f(x)$	16.4	16.94	16.994	17	17.006	17.06	17.6

2. $f(x) = \frac{x - 2}{3x^2 - 4x - 4}$

$\lim_{x \rightarrow 2} f(x) = \frac{1}{8}$

The limit cannot be reached.

x	1.9	1.99	1.999	2	2.001	2.01	2.1
$f(x)$	0.1299	0.1255	0.1250	Undef.	0.1250	0.1245	0.1205

3. $f(x) = \frac{1 - e^{-x}}{x}$

$\lim_{x \rightarrow 0} \frac{1 - e^{-x}}{x} = 1$

The limit cannot be reached.

x	-0.1	-0.01	-0.001	0	0.001	0.01	0.1
$f(x)$	1.0517	1.0050	1.0005	Error	0.9995	0.9950	0.9516

4. $f(x) = \frac{\ln(1 - x)}{x}$

$\lim_{x \rightarrow 0} \frac{\ln(1 - x)}{x} = -1$

The limit cannot be reached.

x	-0.1	-0.01	-0.001	0	0.001	0.01	0.1
$f(x)$	-0.9531	-0.9950	-0.9995	Error	-1.0005	-1.0050	-1.0536

5. $\lim_{x \rightarrow 1} (3 - x) = 2$

6. Limit does not exist.

7. $\lim_{x \rightarrow 1} \frac{x^2 - 1}{x - 1} = 2$

8. $\lim_{x \rightarrow -1} (2x^2 + 1) = 3$

9. (a) $\lim_{x \rightarrow c} [f(x)]^3 = 4^3 = 64$

10. (a) $\lim_{x \rightarrow c} \sqrt[3]{f(x)} = \sqrt[3]{27} = 3$

(b) $\lim_{x \rightarrow c} [3f(x) - g(x)] = 3(4) - 5 = 7$

(b) $\lim_{x \rightarrow c} \frac{f(x)}{18} = \frac{27}{18} = \frac{3}{2}$

(c) $\lim_{x \rightarrow c} [f(x)g(x)] = (4)(5) = 20$

(c) $\lim_{x \rightarrow c} [f(x)g(x)] = (27)(12) = 324$

(d) $\lim_{x \rightarrow c} \frac{f(x)}{g(x)} = \frac{4}{5}$

(d) $\lim_{x \rightarrow c} [f(x) - 2g(x)] = 27 - 2(12) = 3$

11. $\lim_{x \rightarrow 4} (\frac{1}{2}x + 3) = \frac{1}{2}(4) + 3 = 5$

12. $\lim_{x \rightarrow 3} (5x - 4) = 5(3) - 4 = 11$

13. $\lim_{x \rightarrow 2} (5x - 3)(3x + 5) = (5(2) - 3)(3(2) + 5) = (7)(11) = 77$

14. $\lim_{x \rightarrow -2} (5 - 2x - x^2) = 5 - 2(-2) - (-2)^2 = 5$

15. $\lim_{t \rightarrow 3} \frac{t^2 + 1}{t} = \frac{9 + 1}{3} = \frac{10}{3}$

16. $\lim_{x \rightarrow 2} \frac{3x + 5}{5x - 3} = \frac{3(2) + 5}{5(2) - 3} = \frac{11}{7}$

$$17. \lim_{x \rightarrow -2} \sqrt[3]{4x} = (-8)^{1/3} = -2$$

$$18. \lim_{x \rightarrow -1} \sqrt{5-x} = \sqrt{5-(-1)} = \sqrt{6}$$

$$19. \lim_{x \rightarrow \pi} \sin 3x = \sin 3\pi = 0$$

$$20. \lim_{x \rightarrow 0} \tan x = \tan 0 = 0$$

$$21. \lim_{x \rightarrow -1} 2e^x = 2e^{-1} = \frac{2}{e}$$

$$22. \lim_{x \rightarrow 4} \ln x = \ln 4$$

$$23. \lim_{x \rightarrow -1/2} \arcsin x = \arcsin\left(-\frac{1}{2}\right) = -\frac{\pi}{6}$$

$$24. \lim_{x \rightarrow 0} \arctan x = \arctan 0 = 0$$

$$\begin{aligned} 25. \lim_{t \rightarrow -2} \frac{t+2}{t^2-4} &= \lim_{t \rightarrow -2} \frac{t+2}{(t+2)(t-2)} \\ &= \lim_{t \rightarrow -2} \frac{1}{t-2} = -\frac{1}{4} \end{aligned}$$

$$\begin{aligned} 26. \lim_{t \rightarrow 3} \frac{t^2-9}{t-3} &= \lim_{t \rightarrow 3} \frac{(t-3)(t+3)}{t-3} \\ &= \lim_{t \rightarrow 3} (t+3) = 6 \end{aligned}$$

$$\begin{aligned} 27. \lim_{x \rightarrow 5} \frac{x-5}{x^2+5x-50} &= \lim_{x \rightarrow 5} \frac{x-5}{(x-5)(x+10)} \\ &= \lim_{x \rightarrow 5} \frac{1}{x+10} = \frac{1}{15} \end{aligned}$$

$$\begin{aligned} 28. \lim_{x \rightarrow -1} \frac{x+1}{(x^2-5x-6)} &= \lim_{x \rightarrow -1} \frac{(x+1)}{(x+1)(x-6)} \\ &= \lim_{x \rightarrow -1} \frac{1}{x-6} = -\frac{1}{7} \end{aligned}$$

$$\begin{aligned} 29. \lim_{x \rightarrow -2} \frac{x^2-4}{x^3+8} &= \lim_{x \rightarrow -2} \frac{(x+2)(x-2)}{(x+2)(x^2-2x+4)} \\ &= \lim_{x \rightarrow -2} \frac{x-2}{x^2-2x+4} \\ &= \frac{-4}{12} = -\frac{1}{3} \end{aligned}$$

$$\begin{aligned} 30. \lim_{x \rightarrow 4} \frac{x^3-64}{x^2-16} &= \lim_{x \rightarrow 4} \frac{(x-4)(x^2+4x+16)}{(x-4)(x+4)} \\ &= \lim_{x \rightarrow 4} \frac{x^2+4x+16}{x+4} \\ &= \frac{16+16+16}{8} = 6 \end{aligned}$$

$$\begin{aligned} 31. \lim_{x \rightarrow -1} \frac{1/(x+2) - 1}{x+1} &= \lim_{x \rightarrow -1} \frac{1 - (x+2)}{(x+2)(x+1)} \\ &= \lim_{x \rightarrow -1} \frac{-(x+1)}{(x+2)(x+1)} \\ &= \lim_{x \rightarrow -1} \frac{-1}{(x+2)} = -1 \end{aligned}$$


$$\begin{aligned} 32. \lim_{x \rightarrow 0} \frac{(1/(1+x) - 1)}{x} &= \lim_{x \rightarrow 0} \frac{1 - (1+x)}{x(1+x)} \\ &= \lim_{x \rightarrow 0} \frac{-x}{x(1+x)} = -1 \end{aligned}$$

$$\begin{aligned} 33. \lim_{u \rightarrow 0} \frac{\sqrt{4+u} - 2}{u} &= \lim_{u \rightarrow 0} \frac{\sqrt{4+u} - 2}{u} \cdot \frac{\sqrt{4+u} + 2}{\sqrt{4+u} + 2} \\ &= \lim_{u \rightarrow 0} \frac{(4+u) - 4}{u(\sqrt{4+u} + 2)} \\ &= \lim_{u \rightarrow 0} \frac{1}{\sqrt{4+u} + 2} = \frac{1}{4} \end{aligned}$$

$$\begin{aligned} 34. \lim_{v \rightarrow 0} \frac{\sqrt{v+9} - 3}{v} &= \lim_{v \rightarrow 0} \frac{\sqrt{v+9} - 3}{v} \cdot \frac{\sqrt{v+9} + 3}{\sqrt{v+9} + 3} \\ &= \lim_{v \rightarrow 0} \frac{(v+9) - 9}{v[\sqrt{v+9} + 3]} \\ &= \lim_{v \rightarrow 0} \frac{1}{\sqrt{v+9} + 3} = \frac{1}{6} \end{aligned}$$

$$\begin{aligned}
 35. \lim_{x \rightarrow 5} \frac{\sqrt{x-1}-2}{x-5} &= \lim_{x \rightarrow 5} \frac{\sqrt{x-1}-2}{x-5} \cdot \frac{\sqrt{x-1}+2}{\sqrt{x-1}+2} \\
 &= \lim_{x \rightarrow 5} \frac{(x-1)-4}{(x-5)(\sqrt{x-1}+2)} \\
 &= \lim_{x \rightarrow 5} \frac{1}{\sqrt{x-1}+2} = \frac{1}{2+2} = \frac{1}{4}
 \end{aligned}$$


$$\begin{aligned}
 36. \lim_{x \rightarrow 1} \frac{\sqrt{3}-\sqrt{x+2}}{1-x} &= \lim_{x \rightarrow 1} \frac{\sqrt{3}-\sqrt{x+2}}{1-x} \cdot \frac{\sqrt{3}+\sqrt{x+2}}{\sqrt{3}+\sqrt{x+2}} \\
 &= \lim_{x \rightarrow 1} \frac{3-(x+2)}{(1-x)(\sqrt{3}+\sqrt{x+2})} \\
 &= \lim_{x \rightarrow 1} \frac{1}{\sqrt{3}+\sqrt{x+2}} = \frac{1}{2\sqrt{3}} = \frac{\sqrt{3}}{6}
 \end{aligned}$$



$$\lim_{x \rightarrow 3} \frac{x-3}{x^2-9} = \frac{1}{6}$$

(b)


x	2.9	2.99	3	3.01	3.1
$f(x)$	0.1695	0.1669	Error	0.1664	0.1639



$$\lim_{x \rightarrow 4} \frac{4-x}{16-x^2} = \frac{1}{8}$$

(b)


x	3.99	3.999	4	4.001	4.01
y_1	0.12516	0.12502	Error	0.12498	0.12484



$\lim_{x \rightarrow 0} e^{-2/x}$ does not exist.

(b) Answers will vary.


x	-0.1	-0.01	-0.001	0	0.001	0.01	0.1
y_1	4.85 E 8	7.2 E 86	Error	Error	0	1 E -87	2.1 E -9



$$\lim_{x \rightarrow 0} e^{-4/x^2} = 0$$

(b)


x	-0.01	-0.001	0	0.001	0.01
y_1	0	0	Error	0	0



$$\lim_{x \rightarrow 0} \frac{\sin 4x}{2x} = 2$$

(b)


x	-0.1	-0.01	-0.001	0	0.001	0.01	0.1
y_1	1.9471	1.9995	1.999995	error	1.999995	1.995	1.9471



$$\lim_{x \rightarrow 0} \frac{\tan 2x}{x} = 2$$

(b)

x	-0.01	-0.001	0	0.001	0.01
y_1	2.0003	2	Error	2	2.0003




$$\lim_{x \rightarrow 1^+} \frac{\sqrt{2x+1} - \sqrt{3}}{x-1} \approx 0.577$$

$$\left(\text{Exact value: } \frac{\sqrt{3}}{3} \right)$$

(b)

x	1.1	1.01	1.001	1.0001
$f(x)$	0.5680	0.5764	0.5773	0.5773



$$\lim_{x \rightarrow 1^+} \frac{1 - \sqrt{x}}{x-1} = -\frac{1}{2}$$

(b)


x	1.1	1.01	1.001	1.0001
$f(x)$	-0.4881	-0.4988	-0.4999	-0.5000

45. $f(x) = \frac{|x-3|}{x-3}$

Limit does not exist
because

$$\lim_{x \rightarrow 3^+} f(x) = 1 \text{ and}$$


$$\lim_{x \rightarrow 3^-} f(x) = -1.$$



46. $\lim_{x \rightarrow 8^-} \frac{|8-x|}{8-x} = 1$


$$\lim_{x \rightarrow 8^+} \frac{|8-x|}{8-x} = -1$$

$$\lim_{x \rightarrow 8} \frac{|8-x|}{8-x} \text{ does not exist.}$$




47. $f(x) = \frac{2}{x^2 - 4}$


Limit does not exist.



48. $\lim_{x \rightarrow -3} \frac{1}{x^2 + 9} = \frac{1}{(-3)^2 + 9} = \frac{1}{18}$




49. $\lim_{x \rightarrow 5} \frac{|x - 5|}{x - 5}$ does not exist.




50. $\lim_{x \rightarrow -2^-} \frac{|x + 2|}{x + 2} = -1$

$\lim_{x \rightarrow -2^+} \frac{|x + 2|}{x + 2} = 1$

$\lim_{x \rightarrow -2} \frac{|x + 2|}{x + 2}$ does not exist.




51. $\lim_{x \rightarrow 2} f(x)$ does not exist.



52. $\lim_{x \rightarrow 0^-} f(x) = -4$

$\lim_{x \rightarrow 0^+} f(x) = -6$

$\lim_{x \rightarrow 0} f(x)$ does not exist.




$$\begin{aligned} 53. \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h} &= \lim_{h \rightarrow 0} \frac{3(x+h) - (x+h)^2 - (3x - x^2)}{h} \\ &= \lim_{h \rightarrow 0} \frac{3x + 3h - x^2 - 2xh - h^2 - 3x + x^2}{h} = \lim_{h \rightarrow 0} \frac{3h - 2xh - h^2}{h} \\ &= \lim_{h \rightarrow 0} (3 - 2x - h) = 3 - 2x \end{aligned}$$

$$\begin{aligned} 54. \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h} &= \lim_{h \rightarrow 0} \frac{[(x+h)^2 - 5(x+h) - 2] - [x^2 - 5x - 2]}{h} \\ &= \lim_{h \rightarrow 0} \frac{2xh + h^2 - 5h}{h} \\ &= \lim_{h \rightarrow 0} [2x + h - 5] = 2x - 5 \end{aligned}$$


55. Slope ≈ 2

Answers will vary.


56. Slope = 0



Slope at $(2, f(2))$ is approximately 2.




Slope is $\frac{1}{4}$ at $(2, 2)$.



At $(2, f(2)) = (2, -3)$, the slope is approximately -1.5 .



58. At $(2, f(2)) = (2, 2)$.

Slope = -4



60. At $(2, f(2)) = (2, 3)$.

Slope = $\frac{2}{3}$



At $(2, f(2)) = (2, 1)$, the slope is approximately 1.

$$\begin{aligned} 63. \quad m &= \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h} \\ &= \lim_{h \rightarrow 0} \frac{(x+h)^2 - 4(x+h) - (x^2 - 4x)}{h} \\ &= \lim_{h \rightarrow 0} \frac{x^2 + 2xh + h^2 - 4x - 4h - x^2 + 4x}{h} \\ &= \lim_{h \rightarrow 0} \frac{2xh + h^2 - 4h}{h} \\ &= \lim_{h \rightarrow 0} (2x + h - 4) = 2x - 4 \end{aligned}$$

(a) At $(0, 0)$, $m = 2(0) - 4 = -4$.

(b) At $(5, 5)$, $m = 2(5) - 4 = 6$.

$$\begin{aligned} 64. \quad m &= \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h} \\ &= \lim_{h \rightarrow 0} \frac{(1/4)(x+h)^4 - (1/4)x^4}{h} \\ &= \lim_{h \rightarrow 0} \frac{(1/4)[x^4 + 4x^3h + 6x^2h^2 + 4xh^3 + h^4 - x^4]}{h} \\ &= \lim_{h \rightarrow 0} \frac{1}{4}[4x^3 + 6x^2h + 4xh^2 + h^3] = x^3 \end{aligned}$$

(a) At $(-2, 4)$, $m = (-2)^3 = -8$.

(b) At $(1, \frac{1}{4})$, $m = (1)^3 = 1$.

$$\begin{aligned} 65. \quad m &= \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h} = \lim_{h \rightarrow 0} \frac{\frac{4}{x+h-6} - \frac{4}{x-6}}{h} \\ &= \lim_{h \rightarrow 0} \frac{4(x-6) - 4(x+h-6)}{(x+h-6)(x-6)h} \\ &= \lim_{h \rightarrow 0} \frac{-4h}{(x+h-6)(x-6)h} \\ &= \lim_{h \rightarrow 0} \frac{-4}{(x+h-6)(x-6)} = \frac{-4}{(x-6)^2} \end{aligned}$$

(a) At $(7, 4)$, $m = \frac{-4}{(7-6)^2} = -4$.

(b) At $(8, 2)$, $m = \frac{-4}{(8-6)^2} = -1$.

$$\begin{aligned}
 66. \quad m &= \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h} \\
 &= \lim_{h \rightarrow 0} \frac{\sqrt{x+h} - \sqrt{x}}{h} \cdot \frac{\sqrt{x+h} + \sqrt{x}}{\sqrt{x+h} + \sqrt{x}} \\
 &= \lim_{h \rightarrow 0} \frac{(x+h) - x}{h[\sqrt{x+h} + \sqrt{x}]} \\
 &= \lim_{h \rightarrow 0} \frac{1}{\sqrt{x+h} + \sqrt{x}} = \frac{1}{2\sqrt{x}}
 \end{aligned}$$

$$(a) \text{ At } (1, 1), m = \frac{1}{2\sqrt{1}} = \frac{1}{2}.$$

$$(b) \text{ At } (4, 2), m = \frac{1}{2\sqrt{4}} = \frac{1}{4}.$$

$$67. \quad f'(x) = \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h} = \lim_{h \rightarrow 0} \frac{5-5}{h} = 0$$

$$\begin{aligned}
 68. \quad g'(x) &= \lim_{h \rightarrow 0} \frac{g(x+h) - g(x)}{h} \\
 &= \lim_{h \rightarrow 0} \frac{-3 - (-3)}{h} = \lim_{h \rightarrow 0} \frac{0}{h} = 0
 \end{aligned}$$

$$\begin{aligned}
 69. \quad h'(x) &= \lim_{k \rightarrow 0} \frac{h(x+k) - h(x)}{k} \\
 &= \lim_{k \rightarrow 0} \frac{[5 - \frac{1}{2}(x+k)] - [5 - \frac{1}{2}x]}{k} \\
 &= \lim_{k \rightarrow 0} \frac{-\frac{1}{2}k}{k} = -\frac{1}{2}
 \end{aligned}$$

$$\begin{aligned}
 70. \quad f'(x) &= \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h} \\
 &= \lim_{h \rightarrow 0} \frac{3(x+h) - 3x}{h} = 3
 \end{aligned}$$

$$\begin{aligned}
 71. \quad g'(x) &= \lim_{h \rightarrow 0} \frac{g(x+h) - g(x)}{h} \\
 &= \lim_{h \rightarrow 0} \frac{2(x+h)^2 - 1 - (2x^2 - 1)}{h} \\
 &= \lim_{h \rightarrow 0} \frac{2x^2 + 4xh + 2h^2 - 2x^2}{h} \\
 &= \lim_{h \rightarrow 0} (4x + 2h) \\
 &= 4x
 \end{aligned}$$

$$\begin{aligned}
 72. \quad f'(x) &= \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h} = \lim_{h \rightarrow 0} \frac{-(x+h)^3 + 4(x+h) - (-x^3 + 4x)}{h} \\
 &= \lim_{h \rightarrow 0} \frac{-x^3 - 3x^2h - 3xh^2 - h^3 + 4x + 4h + x^3 - 4x}{h} \\
 &= \lim_{h \rightarrow 0} \frac{-3x^2h - 3xh^2 - h^3 + 4h}{h} \\
 &= \lim_{h \rightarrow 0} (-3x^2 - 3xh - h^2 + 4) \\
 &= -3x^2 + 4
 \end{aligned}$$

$$\begin{aligned}
 73. f'(t) &= \lim_{h \rightarrow 0} \frac{f(t+h) - f(t)}{h} \\
 &= \lim_{h \rightarrow 0} \frac{\sqrt{t+h+5} - \sqrt{t+5}}{h} \cdot \frac{\sqrt{t+h+5} + \sqrt{t+5}}{\sqrt{t+h+5} + \sqrt{t+5}} \\
 &= \lim_{h \rightarrow 0} \frac{(t+h+5) - (t+5)}{h(\sqrt{t+h+5} + \sqrt{t+5})} \\
 &= \lim_{h \rightarrow 0} \frac{1}{\sqrt{t+h+5} + \sqrt{t+5}} \\
 &= \frac{1}{2\sqrt{t+5}}
 \end{aligned}$$

$$\begin{aligned}
 74. g'(t) &= \lim_{h \rightarrow 0} \frac{g(t+h) - g(t)}{h} \\
 &= \lim_{h \rightarrow 0} \frac{\sqrt{t+h-3} - \sqrt{t-3}}{h} \cdot \frac{\sqrt{t+h-3} + \sqrt{t-3}}{\sqrt{t+h-3} + \sqrt{t-3}} \\
 &= \lim_{h \rightarrow 0} \frac{(t+h-3) - (t-3)}{h(\sqrt{t+h-3} + \sqrt{t-3})} \\
 &= \lim_{h \rightarrow 0} \frac{1}{\sqrt{t+h-3} + \sqrt{t-3}} \\
 &= \frac{1}{2\sqrt{t-3}}
 \end{aligned}$$

$$\begin{aligned}
 75. g'(s) &= \frac{g(s+h) - g(s)}{h} \\
 &= \lim_{h \rightarrow 0} \frac{\frac{4}{s+h+5} - \frac{4}{s+5}}{h} \\
 &= \lim_{h \rightarrow 0} \frac{4s+20 - 4s - 4h - 20}{(s+h+5)(s+5)h} \\
 &= \lim_{h \rightarrow 0} \frac{-4h}{(s+h+5)(s+5)h} \\
 &= \lim_{h \rightarrow 0} \frac{-4}{(s+h+5)(s+5)} \\
 &= \frac{-4}{(s+5)^2}
 \end{aligned}$$

$$\begin{aligned}
 76. g'(t) &= \lim_{h \rightarrow 0} \frac{g(t+h) - g(t)}{h} \\
 &= \lim_{h \rightarrow 0} \frac{\frac{6}{5-(t+h)} - \frac{6}{5-t}}{h} \\
 &= \lim_{h \rightarrow 0} \frac{30 - 6t - 30 + 6t + 6h}{h(5-t-h)(5-t)} \\
 &= \lim_{h \rightarrow 0} \frac{6}{(5-t-h)(5-t)} \\
 &= \frac{6}{(5-t)^2}
 \end{aligned}$$

$$\begin{aligned}
77. g'(x) &= \lim_{h \rightarrow 0} \frac{g(x+h) - g(x)}{h} \\
&= \lim_{h \rightarrow 0} \frac{\frac{1}{\sqrt{x+h+4}} - \frac{1}{\sqrt{x+4}}}{h} \\
&= \lim_{h \rightarrow 0} \frac{\sqrt{x+4} - \sqrt{x+h+4}}{h\sqrt{x+h+4}\sqrt{x+4}} \cdot \frac{\sqrt{x+4} + \sqrt{x+h+4}}{\sqrt{x+4} + \sqrt{x+h+4}} \\
&= \lim_{h \rightarrow 0} \frac{(x+4) - (x+h+4)}{h\sqrt{x+h+4}\sqrt{x+4}[\sqrt{x+4} + \sqrt{x+h+4}]} \\
&= \lim_{h \rightarrow 0} \frac{-1}{\sqrt{x+h+4}\sqrt{x+4}[\sqrt{x+4} + \sqrt{x+h+4}]} \\
&= \frac{-1}{(x+4)2\sqrt{x+4}} \\
&= \frac{-1}{2(x+4)^{3/2}}
\end{aligned}$$

$$\begin{aligned}
78. f'(x) &= \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h} = \lim_{h \rightarrow 0} \frac{\frac{1}{\sqrt{12-x-h}} - \frac{1}{\sqrt{12-x}}}{h} \\
&= \lim_{h \rightarrow 0} \frac{\sqrt{12-x} - \sqrt{12-x-h}}{h[\sqrt{12-x-h}\sqrt{12-x}]} \cdot \frac{\sqrt{12-x} + \sqrt{12-x-h}}{\sqrt{12-x} + \sqrt{12-x-h}} \\
&= \lim_{h \rightarrow 0} \frac{(12-x) - (12-x-h)}{h[\sqrt{12-x-h}\sqrt{12-x}]} \cdot \frac{1}{[\sqrt{12-x} + \sqrt{12-x-h}]} \\
&= \lim_{h \rightarrow 0} \frac{1}{[\sqrt{12-x-h}\sqrt{12-x}][\sqrt{12-x} + \sqrt{12-x-h}]} \\
&= \frac{1}{(12-x)2\sqrt{12-x}} \\
&= \frac{1}{2(12-x)^{3/2}}
\end{aligned}$$

$$79. \lim_{x \rightarrow \infty} \frac{4x}{2x-3} = \frac{4}{2} = 2$$

$$80. \lim_{x \rightarrow \infty} \frac{7x}{14x+2} = \frac{7}{14} = \frac{1}{2}$$

$$81. \lim_{x \rightarrow -\infty} \frac{2x}{x^2-25} = 0$$

$$82. \lim_{x \rightarrow -\infty} \frac{3x}{(1-x)^3} = 0$$

$$83. \lim_{x \rightarrow \infty} \frac{x^2}{2x+3} \text{ does not exist.}$$

$$84. \lim_{y \rightarrow \infty} \frac{3y^4}{y^2+1} \text{ does not exist.}$$

$$85. \lim_{x \rightarrow \infty} \left[\frac{x}{(x-2)^2} + 3 \right] = 0 + 3 = 3$$

$$86. \lim_{x \rightarrow \infty} \left[2 - \frac{2x^2}{(x+1)^2} \right] = 2 - 2 = 0$$

$$87. a_n = \frac{2n-3}{5n+4}$$

$$a_1 = -\frac{1}{9} \quad a_4 = \frac{5}{24}$$

$$a_2 = \frac{1}{14} \quad a_5 = \frac{7}{29}$$

$$a_3 = \frac{3}{19}$$

$$\lim_{n \rightarrow \infty} a_n = \frac{2}{5}$$

$$88. a_n = \frac{2n}{n^2+1}$$

$$a_1 = \frac{2}{2} = 1 \quad a_4 = \frac{8}{17}$$

$$a_2 = \frac{4}{5} \quad a_5 = \frac{10}{26} = \frac{5}{13}$$

$$a_3 = \frac{6}{10} = \frac{3}{5}$$

$$\lim_{n \rightarrow \infty} a_n = 0$$

$$89. a_n = \frac{(-1)^n}{n^3}$$

$$a_1 = -1 \quad a_4 = \frac{1}{64}$$

$$a_2 = \frac{1}{8} \quad a_5 = -\frac{1}{125}$$

$$a_3 = -\frac{1}{27}$$

$$\lim_{n \rightarrow \infty} a_n = 0$$

$$90. a_n = \frac{(-1)^{n+1}}{n}$$

$$a_1 = 1 \quad a_4 = -\frac{1}{4}$$

$$a_2 = -\frac{1}{2} \quad a_5 = \frac{1}{5}$$

$$a_3 = \frac{1}{3}$$

$$\lim_{n \rightarrow \infty} a_n = 0$$

$$91. a_n = \frac{1}{2n^2}[3 - 2n(n+1)] = \frac{3}{2n^2} - \frac{n+1}{n}$$

$$-0.5, -1.125, -1.16\bar{6}, -1.15625, -1.14$$

$$\lim_{n \rightarrow \infty} a_n = 0 - 1 = -1$$

$$92. a_n = 2 + \frac{2}{n}(n-1) - \frac{4}{n} = 4 - \frac{6}{n}$$

$$-2, 1, 2, \frac{5}{2}, \frac{14}{5}$$

$$\lim_{n \rightarrow \infty} a_n = \lim_{n \rightarrow \infty} \left(4 - \frac{6}{n}\right) = 4$$

$$\begin{aligned} 93. (a) \sum_{i=1}^n \left(\frac{4i^2}{n^2} - \frac{i}{n} \right) \frac{1}{n} &= \frac{4}{n^3} \sum_{i=1}^n i^2 - \frac{1}{n^2} \sum_{i=1}^n i \\ &= \frac{4}{n^3} \frac{n(n+1)(2n+1)}{6} - \frac{1}{n^2} \frac{n(n+1)}{2} \\ &= \frac{4n(n+1)(2n+1) - 3n^2(n+1)}{6n^3} \\ &= \frac{n(n+1)(8n+4-3n)}{6n^3} \\ &= \frac{(n+1)(5n+4)}{6n^2} \end{aligned}$$

$$(c) \lim_{n \rightarrow \infty} S(n) = \frac{5}{6}$$

n	10^0	10^1	10^2	10^3	10^4
$S(n)$	3	0.99	0.8484	0.8348	0.8335

$$\begin{aligned}
 94. \text{ (a) } \sum_{i=1}^n \left[4 - \left(\frac{3i}{n} \right)^2 \right] \left(\frac{3i}{n^2} \right) &= \frac{12}{n^2} \sum_{i=1}^n i - \frac{27}{n^4} \sum_{i=1}^n i^3 = \frac{12}{n^2} \frac{n(n+1)}{2} - \frac{27}{n^4} \frac{n^2(n+1)^2}{4} \\
 &= \frac{24n^2 + 24n - 27(n^2 + 2n + 1)}{4n^2} \\
 &= \frac{-3n^2 - 30n - 27}{4n^2} = \frac{-3}{4n^2} (n^2 + 10n + 9) \\
 &= \frac{-3(n+1)(n+9)}{4n^2}
 \end{aligned}$$

n	10^0	10^1	10^2	10^3	10^4
$S(n)$	-15	-1.5675	-0.8257	-0.7575	-0.7508

(c) $\lim_{n \rightarrow \infty} S_n = -\frac{3}{4}$

$$\begin{aligned}
 95. \text{ Area} &\approx \frac{1}{2} \left(\frac{7}{2} + 3 + \frac{5}{2} + 2 + \frac{3}{2} + 1 \right) \\
 &= \frac{1}{2} \left(\frac{27}{2} \right) = \frac{27}{4} = 6.75
 \end{aligned}$$

$$96. \text{ Width of rectangle: } \frac{1}{4}$$

Height is f evaluated at right endpoint

$$\begin{aligned}
 \text{Area} &\approx \frac{1}{4} \left[f\left(\frac{1}{4}\right) + f\left(\frac{1}{2}\right) + f\left(\frac{3}{4}\right) + f(1) \right] \\
 &= \frac{1}{4} \left[4 - \left(\frac{1}{4}\right)^2 + 4 - \left(\frac{1}{2}\right)^2 + 4 - \left(\frac{3}{4}\right)^2 + 4 - 1 \right] \\
 &= \frac{1}{4} \left[15 - \frac{14}{16} \right] = \frac{113}{32} = 3.53125
 \end{aligned}$$

$$97. f(x) = \frac{1}{4}x^2, \quad b - a = 4 - 0 = 4$$

$$\begin{aligned}
 A &\approx \sum_{i=1}^n f\left(\frac{4i}{n}\right) \left(\frac{4}{n}\right) \\
 &= \sum_{i=1}^n \frac{1}{4} \left(\frac{4i}{n}\right)^2 \left(\frac{4}{n}\right) \\
 &= \frac{1}{n} \sum_{i=1}^n \frac{16}{n^2} i^2 \\
 &= \frac{16}{n^3} \frac{n(n+1)(2n+1)}{6} \\
 &= \frac{8(n+1)(2n+1)}{3n^2}
 \end{aligned}$$

n	4	8	20	50
Approximate area	7.5	6.375	5.74	5.4944

(Exact area is $\frac{16}{3} \approx 5.33$.)

$$98. f(x) = 4x - x^2$$

n	4	8	20	50
Approximate area	10	10.5	10.64	10.6624

(Exact area is $10\frac{2}{3}$.)

$$\begin{aligned}
 99. A &= \lim_{n \rightarrow \infty} \sum_{i=1}^n \left(10 - \frac{10i}{n} \right) \left(\frac{10}{n} \right) \\
 &= \lim_{n \rightarrow \infty} \left[\frac{100}{n} \sum_{i=1}^n 1 - \frac{100}{n^2} \sum_{i=1}^n i \right] \\
 &= \lim_{n \rightarrow \infty} \left[\frac{100}{n} (n) - \frac{100}{n^2} \left(\frac{n(n+1)}{2} \right) \right] \\
 &= \lim_{n \rightarrow \infty} \left[100 - 50 \frac{n(n+1)}{n^2} \right] \\
 &= 100 - 50 = 50, \text{ exact area}
 \end{aligned}$$

$$\begin{aligned}
 100. A &= \lim_{n \rightarrow \infty} \sum_{i=1}^n \left[2 \left(3 + \frac{3i}{n} \right) - 6 \right] \left(\frac{3}{n} \right) \\
 &= \lim_{n \rightarrow \infty} \sum_{i=1}^n \frac{18i}{n^2} = \lim_{n \rightarrow \infty} \frac{18}{n^2} \sum_{i=1}^n i \\
 &= \lim_{n \rightarrow \infty} \frac{18}{n^2} \frac{n(n+1)}{2} = 9, \text{ exact area}
 \end{aligned}$$

$$\begin{aligned}
 101. A &= \lim_{n \rightarrow \infty} \sum_{i=1}^n \left[\left(-1 + \frac{3i}{n} \right)^2 + 4 \right] \left(\frac{3}{n} \right) \\
 &= \lim_{n \rightarrow \infty} \sum_{i=1}^n \left[5 - \frac{6i}{n} + \frac{9i^2}{n^2} \right] \frac{3}{n} \\
 &= \lim_{n \rightarrow \infty} \left[\frac{15}{n} \sum_{i=1}^n 1 - \frac{18}{n^2} \sum_{i=1}^n i + \frac{27}{n^3} \sum_{i=1}^n i^2 \right] \\
 &= \lim_{n \rightarrow \infty} \left[\frac{15}{n}(n) - \frac{18}{n^2} \frac{n(n+1)}{2} + \frac{27}{n^3} \frac{n(n+1)(2n+1)}{6} \right] \\
 &= 15 - 9 + 9 = 15, \text{ exact area}
 \end{aligned}$$

$$\begin{aligned}
 102. A &= \lim_{n \rightarrow \infty} \sum_{i=1}^n 8 \left[\left(\frac{i}{n} \right) - \left(\frac{i}{n} \right)^2 \right] \frac{1}{n} \\
 &= \lim_{n \rightarrow \infty} \left[\frac{8}{n^2} \sum_{i=1}^n i - \frac{8}{n^3} \sum_{i=1}^n i^2 \right] \\
 &= \lim_{n \rightarrow \infty} \left[\frac{8}{n^2} \frac{n(n+1)}{2} - \frac{8}{n^3} \frac{n(n+1)(2n+1)}{6} \right] \\
 &= 4 - \frac{8}{3} = \frac{4}{3}, \text{ exact area}
 \end{aligned}$$

$$103. f(x) = x^3 + 1, [0, 2]$$

The width of each rectangle is $2/n$. The height is

$$f\left(\frac{2i}{n}\right) = \left(\frac{2i}{n}\right)^3 + 1.$$

$$\begin{aligned}
 A &\approx \sum_{i=1}^n \left[\left(\frac{2i}{n} \right)^3 + 1 \right] \left(\frac{2}{n} \right) \\
 &= \sum_{i=1}^n \left[\frac{16i^3}{n^4} + \frac{2}{n} \right] \\
 &= \left(\frac{16}{n^4} \right) \frac{n^2(n+1)^2}{4} + \left(\frac{2}{n} \right) n \\
 &= \frac{4}{n^2} (n+1)^2 + 2
 \end{aligned}$$

$$A = \lim_{n \rightarrow \infty} \left[\frac{4}{n^2} (n+1)^2 + 2 \right] = 4 + 2 = 6$$

$$104. f(x) = 1 - x^3, [-3, -1]$$

The width of each rectangle is $2/n$. The height is

$$\begin{aligned}
 f\left(-3 + \frac{2i}{n}\right) &= 1 - \left(-3 + \frac{2i}{n}\right)^3 \\
 &= 28 - 54\frac{i}{n} + \frac{36i^2}{n^2} - \frac{8i^3}{n^3}.
 \end{aligned}$$

$$\begin{aligned}
 A &\approx \sum_{i=1}^n \left[28 - \frac{54}{n}i + \frac{36}{n^2}i^2 - \frac{8}{n^3}i^3 \right] \left(\frac{2}{n} \right) \\
 &= \frac{56}{n}(n) - \frac{108}{n^2} \frac{n(n+1)}{2} + \frac{72}{n^3} \frac{n(n+1)(2n+1)}{6} - \frac{16}{n^4} \frac{n^2(n+1)^2}{4}
 \end{aligned}$$

$$\begin{aligned}
 A &= \lim_{n \rightarrow \infty} \left[56 - 54 \frac{n(n+1)}{n^2} + 12 \frac{n(n+1)(2n+1)}{n^3} - \frac{4n^2(n+1)^2}{n^4} \right] \\
 &= 56 - 54 + 24 - 4 = 22
 \end{aligned}$$

105. $f(x) = 3(x^3 - x^2)$, $[1, 3]$

The width of each rectangle is $2/n$. The height is

$$f\left(1 + \frac{2i}{n}\right) = 3\left[\left(1 + \frac{2i}{n}\right)^3 - \left(1 + \frac{2i}{n}\right)^2\right]$$

$$= \frac{6i}{n} + \frac{24i^2}{n^2} + \frac{24i^3}{n^3}.$$

$$A \approx \sum_{i=1}^n \left[\frac{6i}{n} + \frac{24i^2}{n^2} + \frac{24i^3}{n^3}\right] \left(\frac{2}{n}\right)$$

$$= \frac{12}{n^2} \frac{n(n+1)}{2} + \frac{48}{n^3} \frac{n(n+1)(2n+1)}{6} + \frac{48}{n^4} \frac{n^2(n+1)^2}{4}$$

$$A = \lim_{n \rightarrow \infty} \left[6 + 8 \frac{n(n+1)(2n+1)}{n^3} + 12 \frac{n^2(n+1)^2}{n^4}\right]$$

$$= 6 + 16 + 12 = 34$$

106. $f(x) = 5 - (x + 2)^2$, $[-2, 0]$

The width of each rectangle is $2/n$. The height is

$$f\left(-2 + \frac{2i}{n}\right) = 5 - \left[\left(-2 + \frac{2i}{n}\right) + 2\right]^2$$

$$= 5 - \left(\frac{2i}{n}\right)^2$$


$$= 5 - \frac{4i^2}{n^2}.$$

$$A \approx \sum_{i=1}^n \left[5 - \frac{4i^2}{n^2}\right] \left(\frac{2}{n}\right) = \sum_{i=1}^n \left[\frac{10}{n} - \frac{8}{n^3} i^2\right]$$

$$A = \lim_{n \rightarrow \infty} \left[\frac{10}{n}(n) - \frac{8}{n^3} \frac{n(n+1)(2n+1)}{6}\right] = 10 - \frac{8}{3} = \frac{22}{3}$$

107. (a) $y = -3.376 \times 10^{-7}x^3 + 3.753 \times 10^{-4}x^2 - 0.168x + 132.168$

(c) Area $\approx 88,868$ square feet; answers will vary.



108. True (assuming all the limits exist)

109. False. The limit does not exist.

110. Answers will vary.

Chapter 11 Practice Test

1. Use a graphing utility to complete the table and use the result to estimate the limit

$$\lim_{x \rightarrow 3} \frac{x - 3}{x^2 - 9}$$

x	2.9	2.99	3	3.01	3.1
$f(x)$?		

2. Graph the function

$$f(x) = \frac{\sqrt{x+4} - 2}{x}$$

and estimate the limit

$$\lim_{x \rightarrow 0} \frac{\sqrt{x+4} - 2}{x}$$

3. Find the limit $\lim_{x \rightarrow 2} e^{x-2}$ by direct substitution.

4. Find the limit $\lim_{x \rightarrow 1} \frac{x^3 - 1}{x - 1}$ analytically.

5. Use a graphing utility to estimate the limit

$$\lim_{x \rightarrow 0} \frac{\sin 5x}{2x}$$

6. Find the limit

$$\lim_{x \rightarrow -2} \frac{|x + 2|}{x + 2}$$

7. Use the limit process to find the slope of the graph of $f(x) = \sqrt{x}$ at the point (4, 2).

8. Find the derivative of the function $f(x) = 3x - 1$.

9. Find the limits.

(a) $\lim_{x \rightarrow \infty} \frac{3}{x^4}$

(b) $\lim_{x \rightarrow -\infty} \frac{x^2}{x^2 + 3}$

(c) $\lim_{x \rightarrow \infty} \frac{|x|}{1 - x}$

10. Write the first four terms of the sequence $a_n = \frac{1 - n^2}{2n^2 + 1}$ and find the limit of the sequence.

11. Find the sum $\sum_{i=1}^{25} (i^2 + i)$.

12. Write the sum $\sum_{i=1}^n \frac{i^2}{n^3}$ as a rational function $S(n)$, and find $\lim_{n \rightarrow \infty} S(n)$.

13. Find the area of the region bounded by $f(x) = 1 - x^2$ over the interval $0 \leq x \leq 1$.